

**BILE-ACID CONJUGATES FOR
PROVIDING SUSTAINED SYSTEMIC CONCENTRATIONS OF DRUGS**

5 **CROSS REFERENCE TO RELATED APPLICATIONS**

This application claims the benefit of U.S. Provisional Application Serial Numbers 60/238,758, filed October 6, 2000, 60/249,804, filed November 17, 2000, and 60/297,472, filed June 11, 2001, all of which are incorporated by reference in their entirety.

10 **BACKGROUND OF THE INVENTION**

Field of the Invention

This invention is directed to compounds that provide for sustained systemic concentrations of therapeutic or prophylactic agents following administration to animals. This invention is also directed to pharmaceutical compositions including and methods using such compounds.

State of the Art

Rapid clearance of drugs from the systemic circulation represents a major impediment to effective clinical use of therapeutic and/or prophylactic compounds. Although multiple factors can influence the systemic concentrations of drugs achieved following administration (including drug solubility, dissolution rate, first-pass metabolism, p-glycoprotein and related efflux mechanisms, hepatic/renal elimination, etc), rapid systemic clearance is a particularly significant reason for suboptimal systemic exposure to many compounds. Rapid systemic clearance may require that large doses of drug be administered to achieve a therapeutic or prophylactic effect. Such larger doses of the drug, however, may result in greater variability in drug exposure, more

frequent occurrence of side effects, or decrease in patient compliance. Frequent drug administration may also be required to maintain systemic drug levels above a minimum effective concentration. This problem is particularly significant for drugs that must be maintained in a well defined concentration window to provide continuous therapeutic or prophylactic benefit while minimizing adverse effects (including for example, antibacterial agents, antiviral agents, anticancer agents, anticonvulsants, anticoagulants, etc.).

Conventional approaches to extend the systemic exposure of drugs with rapid clearance involve the use of formulation or device approaches that provide a slow or sustained release of drug within the intestinal lumen. These approaches are well known in the art and normally require that the drug be well absorbed from the large intestine, where such formulations are most likely to reside while releasing the drug. Drugs that are amenable to conventional sustained release approaches must be orally absorbed in the intestine and traverse this epithelial barrier by passive diffusion across the apical and basolateral membranes of the intestinal epithelial cells. The physicochemical features of a molecule that favor its passive uptake from the intestinal lumen into the systemic circulation include low molecular weight (e.g., < 500 Da), adequate solubility, and a balance of hydrophobic and hydrophilic character (logP generally 1.5-4.0) (Navia and Chaturvedi, 1996).

Polar or hydrophilic compounds are typically poorly absorbed through an animal's intestine as there is a substantial energetic penalty for passage of such compounds across the lipid bilayers that constitute cellular membranes. Many nutrients that result from the digestion of ingested foodstuffs in animals, such as amino acids, di- and tripeptides, monosaccharides, nucleosides and water-soluble vitamins, are polar compounds whose uptake is essential to the viability of the animal. For these substances there exist specific mechanisms for active

transport of the solute molecules across the apical membrane of the intestinal epithelia. This transport is frequently energized by co-transport of ions down a concentration gradient. Solute transporter proteins are generally single sub-unit, multi-transmembrane spanning polypeptides, and upon binding of their
5 substrates are believed to undergo conformational changes which result in movement of the substrate(s) across the membrane.

Over the past 10-15 years, it has been found that a number of orally administered drugs are recognized as substrates by some of these transporter proteins, and that this active transport may largely account for the oral
10 absorption of these molecules (Tsuji and Tamai, 1996). While in most instances the transporter substrate properties of these drugs were unanticipated discoveries made through retrospective analysis, it has been appreciated that, in principle, one might achieve good intestinal permeability for a drug by designing in recognition and uptake by a nutrient transport system. Drugs subject to active
15 absorption in the small intestine are often unable to passively diffuse across epithelial cell membranes and are too large to pass through the tight junctions that exist between the intestinal cells. These drugs include many compounds structurally related to amino acids, dipeptides, sugars, nucleosides, etc. (for example, many cephalosporins, ACE inhibitors, AZT, gabapentin, pregabalin,
20 baclofen, etc.)

Rapid systemic clearance occurs in a portion of these drugs and such clearance requires that they be dosed frequently to maintain a therapeutic or prophylactic concentration in the systemic circulation (Bryans and Wustrow, 1999). Conventional sustained release approaches have not always been
25 successfully applied to these drugs if they are not absorbed from the large intestine. Thus there is a significant need for effective sustained release versions of these drugs, particularly for the pediatric patient population, since drug must

be administered during school hours, raising the issues of compliance, liability, and social acceptance.

One attractive pathway that might be exploitable for sustained delivery of drugs with rapid systemic clearance is the entero-hepatic circulation of bile acids (Swaan *et al*, 1996). Bile acids are hydroxylated steroids that play a key role in digestion and absorption of fat and lipophilic vitamins. After synthesis in the liver, they are secreted into bile and excreted by the gall bladder into the intestinal lumen where they emulsify and help solubilize lipophilic substances. Bile acids are conserved in the body by active uptake from the terminal ileum via the sodium-dependent transporter IBAT (or ASBT) and subsequent hepatic extraction by the transporter NTCP (or LBAT) located in the sinusoidal membrane of hepatocytes. This efficient mechanism to preserve the bile acid pool is termed the enterohepatic circulation (see Figure 2). In man, the total bile acid pool (3-5 g) recirculates 6-10 times per day giving rise to a daily uptake of approximately 20-30 g of bile acids.

The high transport capacity of the bile acid pathway has been a key reason for interest in this system for drug delivery purposes. Several papers have postulated that chemical conjugates of bile acids with drugs could be used to provide liver site-directed delivery of a drug to bring about high therapeutic concentrations in the diseased liver with minimization of general toxic reactions elsewhere in the body; and gallbladder-site delivery systems of cholecystographic agents and cholesterol gallstone dissolution accelerators” (Ho, 1987). Several groups have explored these concepts in some detail, using the C-24 carboxylic acid, C-3, C-7, and C-12 hydroxyl groups of cholic acid (and other bile acids) as handles for chemically conjugating drugs or drug surrogates. (Kramer *et al*, 1992; Kim *et al*, 1993).

5 The most rigorous drug targeting studies using the bile acid transport
pathway to date relate to work with bile acid conjugates of HMG-CoA reductase
inhibitors (Kramer *et al*, 1994b; Petzinger *et al*, 1995; Kramer and Wess, 1995;
Kramer *et al*, 1997b). Coupling of the HMG-CoA reductase inhibitor HR 780
via an amide linkage to the C-3 position of cholate, taurocholate and
glycocholate afforded substrates for both the ileal and liver bile acid transporter
proteins (Figure 3). Upon oral dosing of rats, the cholate conjugate S 3554 led to
specific inhibition of HMG-CoA reductase in the liver, and in contrast to the
parent compound HR 780, gave significantly reduced inhibition of the enzyme
10 in extra-hepatic organs. Companion studies that looked at the tissue distribution
of radiolabeled drugs two hours after intravenous administration through the
mesenteric vein of rats also showed dramatically lower systemic levels for the
bile acid conjugate relative to the parent. Because inhibition of HMG-CoA
reductase requires the presence of the free carboxylic acid moiety in HR 780 this
15 data was taken to indicate that S 3554 served as a prodrug of HR 780,
undergoing hydrolysis (and other uncharacterized metabolism) in the rat liver.
Interestingly, uptake of S 3554 by liver did not appear to depend on the liver bile
acid transporter NTCP (which prefers taurocholate conjugates), but may instead
have involved another multispecific organic anion transport system on the
20 sinusoidal hepatocyte membrane.


In summary, while the concept of harnessing the intestinal bile acid
uptake pathway to enhance the absorption of poorly absorbed drugs is well
appreciated, the existing art has merely demonstrated that bile acid-drug
conjugates may be effectively trafficked to the liver and generally excreted into
25 the bile, either unchanged or as some type of metabolite. The art gives no
guidance as to how one prepares a composition that exploits the bile acid
transport pathway and simultaneously provides therapeutically meaningful
levels of a drug substance outside of the enterohepatic circulation. Furthermore,

the art does not describe the potential use of the bile acid transport pathway to achieve a circulating reservoir of conjugated drug that is slowly released into the systemic circulation to provide sustained concentrations.

SUMMARY OF THE INVENTION

5 This invention is directed to the surprising discovery that the bile acid transport system can be utilized to provide sustained systemic concentrations of drugs administered to an animal. This invention, therefore, permits sustained therapeutic or prophylactic systemic blood concentrations of drugs which heretofore could not be achieved.

10 Accordingly, in one of its compound aspects, this invention is directed to compounds that provide for sustained therapeutic or prophylactic blood concentrations of a drug or an active metabolite thereof in the systemic circulation of an animal. Such compounds are of formula (I):




wherein:

15 R^1 and R^2 are independently hydrogen or hydroxy;
X is selected from the group consisting of hydroxy and Q^x -G- where:
G is -O-, -C(O)O- or -NH-;

Q^x is a group derived from a linear oligopeptide comprising a first moiety D and further comprising from 1 to 3 amino acids, and wherein said group is cleavable from (I) under physiological conditions;

D is a drug containing at least one carboxylic acid group and at least one moiety selected from the group consisting of a primary amino group, a secondary amino group or a hydroxyl group, with the provisos that the drug is not a GABA analog; L-Dopa, an L-aromatic amino acid decarboxylase inhibitor, a catechol O-methyl transferase inhibitor or derivatives thereof; a naturally occurring α -amino acid or an ester or carboxamide of a naturally occurring α -amino acid; a polypeptide derived from a linear oligopeptide containing at least 3 α -amino acids; an oligonucleotide; a cyclophane derivative, a diethylenetriaminopentaacetate derivative, or paramagnetic ion chelates thereof; 5-de-O-methylsporadicin; a bis-(2-chloroethyl)amine containing nitrogen mustard; an HMG-CoA reductase inhibitor; a proline hydroxylase inhibitor; or a steroid containing the carbon substructures of the following formulae:




Z is selected from the group consisting of:

(i) a substituted alkyl group containing a moiety which is negatively charged at physiological pH, which moiety is selected from the group consisting of $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{H}$, $-\text{P}(\text{O})(\text{OR}^6)(\text{OH})$, $-\text{OP}(\text{O})(\text{OR}^6)(\text{OH})$, $-\text{OSO}_3\text{H}$ and the like, and where R^6 is selected from the group consisting of alkyl, substituted alkyl, aryl and substituted aryl; and

(ii) a group of the formula $-\text{M}-\text{Q}^x$, wherein M is selected from the group consisting of $-\text{CH}_2\text{OC}(\text{O})-$ and $-\text{CH}_2\text{CH}_2\text{C}(\text{O})-$, and wherein Q^x is a group

derived from a linear oligopeptide comprising a first moiety D' and further comprising from 1 to 3 amino acids, and wherein said group is cleavable under physiological conditions;

5 D' is a drug containing at least one carboxylic acid group and at least one moiety selected from the group consisting of a primary amino group, a secondary amino group or a hydroxyl group, with the provisos that the drug is not a GABA analog; L-Dopa, an L-aromatic amino acid decarboxylase inhibitor, a catechol O-methyl transferase inhibitor or derivatives thereof; a naturally occurring α -amino acid or an ester or carboxamide of a naturally occurring α -
10 amino acid; a polypeptide derived from a linear oligopeptide containing at least 3 α -amino acids; an oligonucleotide; a cyclophane derivative, a diethylenetriaminopentaacetate derivative, or paramagnetic ion chelates thereof; 5-de-O-methylsporadicin; a bis-(2-chloroethyl)amine containing nitrogen mustard; an HMG-CoA reductase inhibitor; a proline hydroxylase inhibitor; or a
15 steroid containing the carbon substructures of the following formulae:




or a pharmaceutically acceptable salt thereof;

provided that when X is hydroxy, then Z is a group of formula -M-Q^x.

Preferably, when X is Q^x-G-, Q^x is one of the following two structures:





wherein

I is $-\text{NR}^{50}-(\text{CR}^{51}\text{R}^{52})_a-(\text{CR}^{53}\text{R}^{54})_b-\text{C}(\text{O})-$;

J is $-\text{NR}^{55}-(\text{CR}^{56}\text{R}^{57})_c-(\text{CR}^{58}\text{R}^{59})_d-\text{C}(\text{O})-$;

5 K is $-\text{NR}^{60}-(\text{CR}^{61}\text{R}^{62})_e-(\text{CR}^{63}\text{R}^{64})_f-\text{C}(\text{O})-$;

T is $-\text{C}(\text{O})-(\text{CR}^{65}\text{R}^{66})_g-(\text{CR}^{67}\text{R}^{68})_h-\text{NR}^{69}-$;

U is $-\text{C}(\text{O})-(\text{CR}^{70}\text{R}^{71})_m-(\text{CR}^{72}\text{R}^{73})_n-\text{NR}^{74}-$;

V is $-\text{C}(\text{O})-(\text{CR}^{75}\text{R}^{76})_o-(\text{CR}^{77}\text{R}^{78})_p-\text{NR}^{79}-$;

R^{40} is $-\text{OH}$ or $-\text{OR}^{17}$;

10 R^{41} is $-\text{H}$, $-\text{C}(\text{O})\text{R}^{17}$, or $-\text{C}(\text{O})\text{OR}^{17}$;

R^{17} is alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

a, b, c, d, e, f, g, h, m, n, o and p are independently 0 or 1, wherein at
15 least one of a and b is 1, at least one of c and d is 1, at least one of e and f is 1, at least one of g and h is 1, at least one of m and n is 1, at least one of o and p is 1;

i, j, k, t, u and v are independently 0 or 1, wherein at least one of i, j and k is 1, and wherein at least one of t, u and v is 1;

R^{50} is hydrogen or R^{50} and R^{51} together with the atoms to which they are
20 attached form a heterocyclyl ring;

R^{51} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{51} and R^{52} together with the atoms to which they are attached form a
25 cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R^{51} and R^{53} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{52} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

5 R^{53} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{53} and R^{54} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

10 R^{54} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

15 R^{55} is hydrogen or R^{55} and R^{56} , together with the atoms to which they are attached form a heterocyclyl ring;

R^{56} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{56} and R^{57} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R^{56} and R^{58} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

20 R^{57} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

25 R^{58} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl

or R⁵⁸ and R⁵⁹ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R⁵⁹ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R⁶⁰ is hydrogen or R⁶⁰ and R⁶¹, together with the atoms to which they are attached form a heterocyclyl ring;

R⁶¹ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁶¹ and R⁶² together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R⁶¹ and R⁶³ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R⁶² is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R⁶³ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁶³ and R⁶⁴ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R⁶⁴ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

5 R⁶⁵ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁶⁵ and R⁶⁶ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R⁶⁵ and R⁶⁷ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

10 R⁶⁶ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

15 R⁶⁷ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁶⁷ and R⁶⁸ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

20 R⁶⁸ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R⁶⁹ is hydrogen or R⁶⁹ and R⁶⁸ together with the atoms to which they are attached form a heterocyclyl ring;

25 R⁷⁰ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁷⁰ and R⁷¹ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R⁷⁰ and R⁷² together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

5 R⁷² is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁷² and R⁷³ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

15 R⁷⁴ is hydrogen or R⁷⁴ and R⁷³ together with the atoms to which they are attached form a heterocyclyl ring;

R⁷⁶ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R⁷⁷ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl


or R⁷⁷ and R⁷⁸ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R⁷⁸ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl; and

R⁷⁹ is hydrogen or R⁷⁹ and R⁷⁸ together with the atoms to which they are attached form a heterocyclyl ring;

wherein the bond between J_j or U_u and D and any amino acid to which it is attached is an amide or ester bond, and where the bond between D and K_k or T_t is an amide bond.

Preferably, where Z is a substituted alkyl group of the formula -M-Q^{x'}, and wherein Q^{x'} is of the following structure:



wherein

I' is $-\text{NR}^{50'}-(\text{CR}^{51'}\text{R}^{52'})_a-(\text{CR}^{53'}\text{R}^{54'})_b-\text{C}(\text{O})-$;

J' is $-\text{NR}^{55'}-(\text{CR}^{56'}\text{R}^{57'})_c-(\text{CR}^{58'}\text{R}^{59'})_d-\text{C}(\text{O})-$;

K' is $-\text{NR}^{60'}-(\text{CR}^{61'}\text{R}^{62'})_e-(\text{CR}^{63'}\text{R}^{64'})_f-\text{C}(\text{O})-$;

R^{40'} is -OH or -OR^{17'};

R^{17'} is alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

a', b', c', d', e' and f' are independently 0 or 1, wherein at least one of a' and b' is 1, at least one of c' and d' is 1, at least one of e' and f' is 1;

i', j' and k' are independently 0 or 1, wherein at least one of i', j' and k' is 1;

$R^{50'}$ is hydrogen or $R^{50'}$ and $R^{51'}$ together with the atoms to which they are attached form a heterocyclyl ring;

$R^{51'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{51'}$ and $R^{52'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or $R^{51'}$ and $R^{53'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

$R^{52'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

$R^{53'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{53'}$ and $R^{54'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

$R^{54'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

$R^{55'}$ is hydrogen or $R^{55'}$ and $R^{56'}$, together with the atoms to which they are attached form a heterocyclyl ring;

$R^{56'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{56'}$ and $R^{57'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring,

or R^{56'} and R^{58'} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{57'} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R^{58'} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{58'} and R^{59'} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{59'} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R^{60'} is hydrogen or R^{60'} and R^{61'}, together with the atoms to which they are attached form a heterocyclyl ring;

R^{61'} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{61'} and R^{62'} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R^{61'} and R^{63'} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{62'} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

106007-89242660

$R^{63'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{63'}$ and $R^{64'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

$R^{64'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

wherein the bond between $J'_{j'}$ and D' and any amino acid to which it is attached is an amide or ester bond.

Preferably, R^1 and R^2 of a compound of formula (I) are both α -OH; R^1 is β -OH and R^2 is hydrogen; R^1 is α -OH and R^2 is hydrogen; R^1 is hydrogen and R^2 is α -OH; R^1 is β -OH and R^2 is α -OH; or R^1 and R^2 are both hydrogen

When amino acids I, J, K, T, U and V are included in a compound of formula (I), they are typically derived from naturally occurring α -amino acids.

When amino acids I' , J' and K' are included in a compound of formula (I), they are typically derived from naturally occurring α -amino acids.


Preferably, where Q^x of a compound of formula (I) is of the structure $-I_i-$
 $J_j-D-K_k-R^{40}$ or $-T_t-D-U_u-V_v-R^{41}$, then b, c, d, e, f, g, h, j, k, m, n, o and p are 0, and a' and i' are 1.

Preferably, where $Q^{x'}$ of a compound of formula (I) is of the structure $-I'_{i'}-J'_{j'}-D'-K'_k-R^{40'}$, then b', c', d', e' and f' are 0, and a' is 1.

Preferably, with respect to substituents included in a compound of formula (I), X is hydroxy and Q^{x'} is -I'_i-J'_j-D'-K'_k-R^{40'}.

Preferably in the method, where a compound of formula (I) contains the substituents R^{50'}, R^{51'} and R^{52'}, the substituents are defined as follows: R^{50'} is hydrogen, R^{51'} is selected from the group consisting of hydrogen, methyl, 2-propyl, 2-butyl, isobutyl, *t*-butyl, cyclopentyl, cyclohexyl, phenyl, benzyl, 4-hydroxybenzyl, 2-imidazolyl, 2-indolyl, -CH₂OH, -CH(OH)CH₃, -CH₂CONH₂, -CH₂CH₂CONH₂, -CH₂CH₂SCH₃ or CH₂SH, and R^{52'} is hydrogen.

Another aspect of this invention is a compound of formula (II):



10 wherein:

R¹ and R² are independently hydrogen or hydroxy;


X is selected from the group consisting of hydroxy and P^x-G- where:

G is -O-, -C(O)O- or -NH-;

15 P^x is a group derived from a linear oligopeptide comprising a first moiety D'' and further comprising from 1 to 3 amino acids, and wherein said group is cleavable from (II) under physiological conditions;

D'' is a drug containing at least one moiety selected from the group consisting of a primary amino group, a secondary amino group or a hydroxyl group, with the provisos that the drug is not a GABA analog; L-Dopa, an L-

aromatic amino acid decarboxylase inhibitor, a catechol O-methyl transferase inhibitor or derivatives thereof; a naturally occurring α -amino acid or an ester or carboxamide of a naturally occurring α -amino acid; a polypeptide derived from a linear oligopeptide containing at least 3 α -amino acids; an oligonucleotide; a
 5 cyclophane derivative, a diethylenetriaminopentaacetate derivative, or paramagnetic ion chelates thereof; histamine or tyramine; 5-de-O-methylsporaricin; a bis-(2-chloroethyl)amine containing nitrogen mustard; an HMG-CoA reductase inhibitor; a proline hydroxylase inhibitor; fluvalinate; or a steroid containing the carbon substructures of the following formulae:




10 Z is selected from the group consisting of:

(i) a substituted alkyl group containing a moiety which is negatively charged at physiological pH, which moiety is selected from the group consisting of $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{H}$, $-\text{P}(\text{O})(\text{OR}^6)(\text{OH})$, $-\text{OP}(\text{O})(\text{OR}^6)(\text{OH})$, $-\text{OSO}_3\text{H}$ and the like, and where R^6 is selected from the group consisting of alkyl, substituted
 15 alkyl, aryl and substituted aryl;
 and

(ii) a group of the formula $-\text{M}-\text{P}^{\text{x}'}$, wherein M is selected from the group consisting of $-\text{CH}_2\text{OC}(\text{O})-$ and $-\text{CH}_2\text{CH}_2\text{C}(\text{O})-$, and wherein $\text{P}^{\text{x}'}$ is a group derived from a linear oligopeptide comprising a first moiety $\text{D}^{\text{'''}}$ and further
 20 comprising from 1 to 3 amino acids, and wherein said group is cleavable under physiological conditions;


$\text{D}^{\text{'''}}$ is a drug containing at least one moiety selected from the group consisting of a primary amino group, a secondary amino group or a hydroxyl

group, with the provisos that the drug is not a GABA analog; L-Dopa, an L-aromatic amino acid decarboxylase inhibitor, a catechol O-methyl transferase inhibitor or derivatives thereof; a naturally occurring α -amino acid or an ester or carboxamide of a naturally occurring α -amino acid; a polypeptide derived from
 5 a linear oligopeptide containing at least 3 α -amino acids; an oligonucleotide; a cyclophane derivative, a diethylenetriaminopentaacetate derivative, or paramagnetic ion chelates thereof; histamine or tyramine; 5-de-O-methylsporaricin; a bis-(2-chloroethyl)amine containing nitrogen mustard; an HMG-CoA reductase inhibitor; a proline hydroxylase inhibitor; fluvalinate; or a
 10 steroid containing the carbon substructures of the following formulae:



or a pharmaceutically acceptable salt thereof;
 provided that when X is hydroxy, then Z is a group of formula -M-P^x.

Preferably, X is P^x-G-, G is -C(O)O-, and P^x is of the following structure:



15 wherein

I is $-\text{NR}^{50}-(\text{CR}^{51}\text{R}^{52})_a-(\text{CR}^{53}\text{R}^{54})_b-\text{C(O)}-;$

J is $-\text{NR}^{55}-(\text{CR}^{56}\text{R}^{57})_c-(\text{CR}^{58}\text{R}^{59})_d-\text{C(O)}-;$

K is $-\text{NR}^{60}-(\text{CR}^{61}\text{R}^{62})_e-(\text{CR}^{63}\text{R}^{64})_f-\text{C(O)}-;$

20 a, b, c, d, e and f are independently 0 or 1, wherein at least one of a and b is 1, at least one of c and d is 1, at least one of e and f is 1;

i, j and k are independently 0 or 1, wherein at least one of i, j and k is 1;

R^{50} is hydrogen or R^{50} and R^{51} together with the atoms to which they are attached form a heterocyclyl ring;

R^{51} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{51} and R^{52} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R^{51} and R^{53} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{52} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R^{53} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{53} and R^{54} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{54} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R^{55} is hydrogen or R^{55} and R^{56} , together with the atoms to which they are attached form a heterocyclyl ring;

R^{56} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{56} and R^{57} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring,

or R⁵⁶ and R⁵⁸ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

5 R⁵⁷ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

10 R⁵⁸ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁵⁸ and R⁵⁹ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

15 R⁵⁹ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R⁶⁰ is hydrogen or R⁶⁰ and R⁶¹, together with the atoms to which they are attached form a heterocyclyl ring;


20 R⁶¹ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁶¹ and R⁶² together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R⁶¹ and R⁶³ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

25 R⁶² is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

5 R^{63} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{63} and R^{64} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

10 R^{64} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;
wherein the bond between K_k and D'' and any amino acid to which it is attached is an amide or ester bond.

Preferably, where Z is a substituted alkyl group of the formula $-M-P^x$, P^x is of the following structure:



15 wherein
 I' is $-[NR^{50'}-(CR^{51'}R^{52'})_a-(CR^{53'}R^{54'})_b-C(O)]-$;
 J' is $-[NR^{55'}-(CR^{56'}R^{57'})_c-(CR^{58'}R^{59'})_d-C(O)]-$;
 K' is $-[NR^{60'}-(CR^{61'}R^{62'})_e-(CR^{63'}R^{64'})_f-C(O)]-$;
 a', b', c', d', e' and f' are independently 0 or 1, wherein at least one of a' and b' is 1, at least one of c' and d' is 1, at least one of e' and f' is 1;
20 i', j' and k' are independently 0 or 1, wherein at least one of i', j' and k' is 1;;

$R^{50'}$ is hydrogen or $R^{50'}$ and $R^{51'}$ together with the atoms to which they are attached form a heterocyclyl ring;
25 $R^{51'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl

or $R^{51'}$ and $R^{52'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or $R^{51'}$ and $R^{53'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

5 $R^{52'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

10 $R^{53'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{53'}$ and $R^{54'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

15 $R^{54'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

$R^{55'}$ is hydrogen or $R^{55'}$ and $R^{56'}$, together with the atoms to which they are attached form a heterocyclyl ring;

20 $R^{56'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{56'}$ and $R^{57'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or $R^{56'}$ and $R^{58'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

25 $R^{57'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl,

substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

5 $R^{58'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{58'}$ and $R^{59'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

10 $R^{59'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

$R^{60'}$ is hydrogen or $R^{60'}$ and $R^{61'}$, together with the atoms to which they are attached form a heterocyclyl ring;

15 $R^{61'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{61'}$ and $R^{62'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or $R^{61'}$ and $R^{63'}$ together with the atoms to which they are attached form a
20 cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

$R^{62'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

25 $R^{63'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or $R^{63'}$ and $R^{64'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

106007-2942660

$R^{64'}$ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

- 5 wherein the bond between K'_k and D''' and any amino acid to which it is attached is an amide or ester bond.

Preferably, R^1 and R^2 are both α -OH; R^1 is β -OH and R^2 is hydrogen; R^1 is α -OH and R^2 is hydrogen; R^1 is hydrogen and R^2 is α -OH; R^1 is β -OH and R^2 is α -OH; or R^1 and R^2 are both hydrogen.

- 10 When amino acids I, J and K are included in a compound of Formula II, they are typically derived from naturally occurring α -amino acids.


When amino acids I' , J' and K' are included in a compound of Formula II, they are typically derived from naturally occurring α -amino acids.

- 15 Preferably, where P^x of a compound of formula II is of the structure $-I_i-J_j-K'_k-D''$ in Formula II, b, c, d, e, f, j and k are 0, and a and i are 1.

Preferably, where P^x of a compound of formula II is of the structure $-I'_{i'}-J'_{j'}-K'_{k'}-D'''$, b', c', d', e', f', j' and k' are 0, and wherein a' and i' are 1.

Yet another aspect of this invention is a compound of formula (III):

Publ. No. 2004/012660



wherein:


R¹ and R² are independently hydrogen or hydroxy;

X is selected from the group consisting of hydroxy and S^x-G- where:

G is -O-, or -NH-;

5 S^x is a group derived from a linear oligopeptide comprising a first moiety D* and further comprising from 1 to 3 amino acids, and wherein said group is cleavable from (III) under physiological conditions;


D* is a drug containing at least one carboxylic acid group, with the provisos that the drug is not a GABA analog; L-Dopa, an L-aromatic amino acid
 10 decarboxylase inhibitor, a catechol O-methyl transferase inhibitor or derivatives thereof; a naturally occurring α-amino acid or an ester or carboxamide of a naturally occurring α-amino acid; a polypeptide derived from a linear oligopeptide containing at least 3 α-amino acids; an oligonucleotide; a cyclophane derivative, a diethylenetriaminopentaacetate derivative, or
 15 paramagnetic ion chelates thereof; 5-de-O-methylsporaricin; a bis-(2-chloroethyl)amine containing nitrogen mustard; an HMG-CoA reductase inhibitor; a proline hydroxylase inhibitor; or a steroid containing the carbon substructures of the following formulae:



Z is selected from the group consisting of:

a substituted alkyl group containing a moiety which is negatively charged at physiological pH, which moiety is selected from the group consisting of $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{H}$, $-\text{P}(\text{O})(\text{OR}^6)(\text{OH})$, $-\text{OP}(\text{O})(\text{OR}^6)(\text{OH})$, $-\text{OSO}_3\text{H}$ and the like, and where R^6 is selected from the group consisting of alkyl, substituted alkyl, aryl and substituted aryl.

Preferably, X is $\text{S}^x\text{-G-}$, and S^x is of the following structure:



wherein:

T is $-\text{[C(O)}-(\text{CR}^{65}\text{R}^{66})_g-(\text{CR}^{67}\text{R}^{68})_h-\text{NR}^{69}]$ -;

U is $-\text{[C(O)}-(\text{CR}^{70}\text{R}^{71})_m-(\text{CR}^{72}\text{R}^{73})_n-\text{NR}^{74}]$ -;

V is $-\text{[C(O)}-(\text{CR}^{75}\text{R}^{76})_o-(\text{CR}^{77}\text{R}^{78})_p-\text{NR}^{79}]$ -;

g, h, m, n, o and p are independently 0 or 1, wherein at least one of g and h is 1, at least one of m and n is 1, at least one of o and p is 1;

t, u and v are independently 0 or 1, wherein at least one of t, u and v is 1;

R^{65} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{65} and R^{66} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring,

or R⁶⁵ and R⁶⁷ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R⁶⁶ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R⁶⁷ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁶⁷ and R⁶⁸ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R⁶⁸ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R⁶⁹ is hydrogen or R⁶⁹ and R⁶⁸ together with the atoms to which they are attached form a heterocyclyl ring;

R⁷⁰ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁷⁰ and R⁷¹ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R⁷⁰ and R⁷² together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R⁷¹ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R^{72} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{72} and R^{73} together with the atoms to which they are attached form a
5 cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{73} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

10 R^{74} is hydrogen or R^{74} and R^{73} together with the atoms to which they are attached form a heterocyclyl ring;

R^{75} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{75} and R^{76} together with the atoms to which they are attached form a
15 cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R^{75} and R^{77} together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{76} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

R^{77} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R^{77} and R^{78} together with the atoms to which they are attached form a
25 cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

R^{78} is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl,

09974768 100007 89242660

substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl; and

R^{79} is hydrogen or R^{79} and R^{78} together with the atoms to which they are attached form a heterocyclyl ring;

5 wherein the bond between V_v and D^* and any amino acid to which it is attached is an amide bond.

Preferably, R^1 and R^2 of Formula III are both α -OH; R^1 is β -OH and R^2 is hydrogen; R^1 is α -OH and R^2 is hydrogen; R^1 is hydrogen and R^2 is α -OH; R^1 is β -OH and R^2 is α -OH; or R^1 and R^2 are both hydrogen.

10 Preferably, when included in a compound of Formula III, amino acids T, U and V are derived from naturally occurring α -amino acids.

Preferably, where S^x of a compound of formula III is of the structure $-T_t-U_u-V_v-D^*$, then h, m, n, o, p, u and v are 0, and g and t are 1.

15 In a method aspect of this invention, the compounds of this invention are preferably used in a method for achieving sustained therapeutic or prophylactic blood concentrations of a drug, or an active metabolite thereof, in the systemic circulation of an animal. The method involves administering a compound of formula (I), (II), or (III) to an animal.

20 In a composition aspect of this invention, the compounds of this invention are mixed with a pharmaceutically acceptable carrier to provide a composition. The composition is preferably used in the method of achieving sustained therapeutic or prophylactic blood concentrations of a drug or an active metabolite thereof discussed above.

105001-8944660

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates structural analogs of γ -aminobutyric acid (GABA).

FIG. 2 illustrates the enterohepatic circulation with key transporter proteins identified which mediate bile acid circulation.

FIG. 3 illustrates HMG-CoA reductase inhibitor HR 780 as well as conjugates employing the lactone-opened ring of HR 780 coupled to a bile acid.

FIGs. 4-8 illustrate reaction sequences for preparation of various compounds having a GABA analog substituent, similar to the compounds of Formula I.

DETAILED DESCRIPTION OF THE INVENTION

This invention is directed to compounds that provide for sustained systemic concentrations of therapeutic or prophylactic drugs following administration to animals. This invention is also directed to methods using the compounds and pharmaceutical compositions that are used in such methods.

However, prior to describing this invention in further detail, the following terms will first be defined:

Definitions

As used herein, the term "animal" refers to various species such as mammalian and avian species including, by way of example, humans, cattle, sheep, horses, dogs, cats, turkeys, chicken, and the like. Preferably, the animal is a mammal and even more preferably is a human.

"Administering to the animal" refers to delivering a compound of formula (I) to an animal through a suitable route. Such routes include, for example, oral, rectal, subcutaneous, intravenous, intramuscular and intranasal. Preferably, the compound is orally administered to the animal.

“Orally delivered drugs” refer to drugs which are administered to an animal in an oral form, preferably, in a pharmaceutically acceptable diluent. Oral delivery includes ingestion of the drug as well as oral gavage of the drug.

5 “PEPT1 oligopeptide transporter” refers to a type of protein that absorbs peptides in certain tissues, such as the intestine. This transporter is described and characterized in the literature. See Adibi, S. A., *Gastroenterology* **1997**, *113*, 332-340 and Leibach *et al.*, *Ann. Rev. Nutr.* **1996**, *16*, 99-119 for a discussion of the transporter.

10 “PEPT2 oligopeptide transporter” refers to a type of protein that absorbs peptides in certain tissues, such as the kidney. This transporter is described and characterized in the literature. See Dieck, S. T. et al., *GLIA* **1999**, *25*, 10-20, Leibach *et al.*, *Ann. Rev. Nutr.* **1996**, *16*, 99-119; and Wong *et al.*, *Am. J. Physiol.* **1998**, *275*, C967-C975 for a discussion of the transporter.

15 “Transported by either a PEPT1 or PEPT2 oligopeptide transporter” refers to the translocation of a molecule across a membrane of a cell expressing the transporter. The translocation occurs through interaction with the transporter and is energized by cotransport of H⁺ ions across the membrane.


20 “Amino acid” is intended to denote α-amino acids and β-amino acids only. α-Amino acids are molecules of the formula $\text{HNR}^{50}\text{-CR}^{51}\text{R}^{52}\text{-C(O)OH}$: wherein:

R⁵⁰ is hydrogen or R⁵⁰ and R⁵¹ together with the atoms to which they are attached form a heterocyclyl ring;

25 R⁵¹ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl

or R⁵¹ and R⁵² together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring.

β-Amino acids are molecules of formula:



5 wherein:

R⁵⁰ is hydrogen or R⁵⁰ and R⁵¹ together with the atoms to which they are attached form a heterocyclyl ring;

10 R⁵¹ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁵¹ and R⁵² together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring, or R⁵¹ and R⁵³ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

15 R⁵² is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

20 R⁵³ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or R⁵³ and R⁵⁴ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclyl or substituted heterocyclyl ring;

25 R⁵⁴ is hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl.

0974758 10094
T06001: 89242660

“Naturally occurring amino acid” refers to any of the alpha-amino acids that are the chief components of proteins. The amino acids are either synthesized by living cells or are obtained as essential components of the diet. Such amino acids include, for example, the following: alanine, arginine, asparagines, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine.

“Drugs that are either completely or incompletely translocated across the intestinal wall into the systemic blood circulation of an animal” refer to any of the well known orally delivered drugs currently delivered by oral administration as well as drugs which cannot be orally administered because such drugs are insufficiently translocated across the intestinal wall of an animal to provide therapeutic or prophylactic blood concentrations in said animal.

Preferably, drugs that fall into the following categories:

- (i) drugs which are insufficiently translocated across the intestinal wall to provide therapeutic or prophylactic blood concentrations;
- (ii) incompletely translocated drugs; or
- (iii) drugs that are either completely or incompletely translocated across the intestinal wall into the systemic blood circulation of an animal, contain suitable functionality to provide points of linkage in forming compounds of formula (I), (II) and (III) above. Such functionality includes, by way of example, carboxyl groups, amine groups and hydroxyl groups.

Examples of drugs containing carboxyl groups include, for instance, angiotensin-converting enzyme inhibitors such as alecapril, 1-[4-carboxy-2-methyl-2R,4R-pentanoyl]-2,3-dihydro-2S-indole-2-carboxylic acid, enalaprilic acid, lisinopril, N-cyclopentyl-N-[3-[(2,2-dimethyl-1-

oxopropyl)thio]-2-methyl-1-oxopropyl]glycine, pivopril, quinaprilat, (2R, 4R)-
2-hydroxyphenyl)-3-(3-mercaptopropionyl)-4-thiazolidinecarboxylic acid, (S)
benzamido-4-oxo-6-phenylhexenoyl-2-carboxypyrrolidine,
[2S-1 [R*(R*))]] 2 α , 3 α β , 7 α β]-1 [2-[[1-carboxy-3-phenylpropyl]-amino]-1-oxopropyl]octahydro-1H-indole-2-carboxylic acid,
5 [3S-1[R*(R*))]]], 3R*]-2-[2-[[1-carboxy-3-phenylpropyl]-amino]-1-oxopropyl]-
1,2,3,4-tetrahydro-3-isoquinolone carboxylic acid and tiopronin; cephalosporin
antibiotics such as cefaclor, cefadroxil, cefamandole, cefatrizine, cefazedone,
cefazuflur, cefazolin, cefbuperazone, cefixime, cefmenoxime, cefmetazole,
10 cefodizime, cefonicid, cefoperazone, ceforanide, cefotaxime, cefotefan,
cefotiam, cefoxitin, cefpimizole, cefpirome, cefpodoxime, cefroxadine,
cefsulodin, cefpiramide, ceftazidime, ceftazole, ceftizoxime, ceftriaxone,
cefuroxime, cephradine, cephaloglycin, cephaloridine,
cephalosporin, cephanone, cephradine and latamoxef; penicillins such as
15 amoxycillin, ampicillin, apalcillin, azidocillin, azlocillin, benzylpenicillin,
carbenicillin, carfecillin, carindacillin, cloxacillin, cyclacillin, dicloxacillin,
epicillin, flucloxacillin, hetacillin, methicillin, mezlocillin, nafcillin, oxacillin,
phenethicillin, piperazillin, sulbenicillin, temocillin and ticarcillin; thrombin
inhibitors such as argatroban, melagatran and napsagatran; influenza
20 neuraminidase inhibitors such as zanamivir and BCX-1812; non-steroidal
antiinflammatory agents such as acetaminophen, alclofenac, alminoprofen, aspirin
(acetylsalicylic acid), 4-biphenylacetic acid, bucloxic acid, carprofen,
cinchophen, cinmetacin, clometacin, clonixin, diclofenac, diflunisal, etodolac,
fenbufen, fenclofenac, fencloxic acid, fenoprofen, ferobufen, flufenamic acid,
25 flufenisal, flurbiprofen, fluprofen, flutiazin, ibufenac, ibuprofen, indomethacin,
indoprofen, ketoprofen, ketorolac, lonazolac, loxoprofen, meclofenamic acid,
mefenamic acid,

1050113924660

2-(8-methyl-10,11-dihydro-11-oxodibenz[b,f]oxepin-2-yl)propionic acid,
naproxen, niflumonic acid, O-(carbamoylphenoxy)acetic acid, oxoprozin,
pirprofen, prodolic acid, salicylic acid, salicylsalicylic acid, sulindac,
suprofen, tiaprofenic acid, tolfenamic acid, tolmetin and zopemirac;
5 prostaglandins such as ciprostone, 16-deoxy-16-hydroxy-16-vinyl prostaglandin
E₂, 6,16-dimethylprostaglandin E₂, epoprostenol, meteneprost, nileprost,
prostacyclin, prostaglandins E₁, E₂, or F_{2α} and thromboxane A₂; quinolone
antibiotics such as acrosoxacin, cinoxacin, ciprofloxacin, enoxacin,
flumequine, naladixic acid, norfloxacin, ofloxacin, oxolinic acid, pefloxacin,
10 pipemidic acid and piromidic acid; other antibiotics such as aztreonam,
imipenem, meropenem and related carbopenem antibiotics.

Representative drugs containing amine groups include: acebutalol,
albuterol, alprenolol, atenolol, bunolol, bupropion, butopamine, butoxamine,
carbuterol, cartelolol, colterol, deterenol, dexpropanolol, diacetolol,
15 dobutamine, exaprolol, exprenolol, fenoterol, fenyrilol, labotolol,
levobunolol, metolol, metaproterenol, metoprolol, nadolol, pamatolol,
penbutalol, pindolol, pirbuterol, practolol, prenalterol, primidolol, prizidilol,
procaterol, propanolol, quinterenol, rimiterol, ritodrine, solotol, soterenol,
sulfiniolol, sulfinterol, sulictidil, tazaolol, terbutaline, timolol, tiprenolol,
20 tipridil, tolamolol, thiabendazole, albendazole, albutoin, alendronate, alinidine,
alizapride, amiloride, aminorex, aprinocid, cambendazole, cimetidine,
cisapride, clonidine, cyclobenzadole, delavirdine, efegatrin, etintidine,
fenbendazole, fenmetazole, flubendazole, fludorex, icadronate, lobendazole,
mebendazole, metazoline, metoclopramide, methylphenidate, mexiletine,
25 neridronate, nocodazole, oxfendazole, oxibendazole, oxmetidine, pamidronate,
parbendazole, pramipexole, prazosin, procainamide, ranitidine,

000001-3924660

tetrahydrazoline, tiamenidine, tinazoline, tiotidine, tocainide, tolazoline, tramazoline, xylometazoline, dimethoxyphenethylamine,

N-[3(R)-[2-piperidin-4-yl)ethyl]-2-piperidone-1-yl]acetyl-3(R)-methyl-
β-alanine, adrenolone, aletamine, amidephrine, amphetamine, aspartame,
5 bamethan, betahistine, clorprenaline, chlortermine, ephrinephrine etryptamine,
fenfluramine, methyldopamine, norepinephrine, tocainide, enviroxime,
nifedipine, nimodipine, triamterene, norfloxacin and similar compounds such
as pipedemic acid,
1-ethyl-6-fluoro-1,4dihydro-4-oxo-7-(1-piperazinyl)-1,8-napthyridine-3-
10 carboxylic acid, 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(piperazinyl)-3-
quinolinecarboxylic acid.

Representative drugs containing hydroxy groups include: steroidal
hormones such as allylestrenol, cingestol, dehydroepiandrosteron, dienostrol,
diethylstilbestrol, dimethisteron, ethyneron, ethynodiol, estradiol, estron,
15 ethinyl estradiol, ethisteron, lynestrenol, mestranol, methyl testosterone,
norethindron, norgestrel, norvinsteron, oxogeston, quinestrol, testosterone and
tigestol; tranquilizers such as dofexazepam, hydroxyzin, lorazepam and
oxazepam; neuroleptics such as acetophenazine, carphenazine, fluphenazine,
perphenyzine and piperazetazine; cytostatics such as aclarubicin, cytarabine,
20 decitabine, daunorubicin, dihydro-5-azacytidine, doxorubicin, epirubicin,
estramustin, etoposide, fludarabine, gemcitabine, 7-hydroxychlorpromazin,
nelarabine, neplanocin A, pentostatin, podophyllotoxin, tezacitabine,
troxacitabine, vinblastin, vincristin, vindesin; hormones and hormone
antagonists such as buserilin, gonadoliberin, icatibrant and leuprorelin acetate;
25 antihistamines such as terphenadine; analgesics such as diflunisal, naproxol,
paracetamol, salicylamide and salicylic acid; antibiotics such as

106001" 894260


azidamphenicol, azithromycin, camptothecin, cefamandol, chloramphenicol, clarithromycin, clavulanic acid, clindamycin, demeclocyclin, doxycyclin, erythromycin, gentamycin, imipenem, latamoxef, metronidazole, neomycin, novobiocin, oleandomycin, oxytetracyclin, tetracycline, thiamenicol and tobramycin; antivirals such as acyclovir, d4C, ddC, DMDC, Fd4C, FddC, FMAU, FTC, 2'-fluoro-ara-dideoxyinosine, ganciclovir, lamivudine, penciclovir, SddC, stavudine, 5-trifluoromethyl-2'-deoxyuridine, zalcitabine and zidovudine; bisphosphonates such as EB-1053, etidronate, ibandronate, olpadronate, residronate, YH-529 and zolendronate; protease inhibitors such as ciprokiren, enalkiren, ritonavir, saquinavir and terlakiren; prostaglandins such as arbaprostil, carboprost, misoprostil and prostacydin; antidepressives such as 8-hydroxychlorimipramine and 2-hydroxyimipramine; antihypertensives such as sotalol and fenoldopam; anticholinergics such as biperidine, procyclidin and trihexyphenidol; antiallergenics such as cromolyn; glucocorticoids such as betamethasone, budenosid, chlorprednisol, clobetasol, clobetasone, corticosteron, cortisone, cortodexon, dexamethason, flucortolon, fludrocortisone, flumethasone, flunisolid, fluprednisolon, flurandrenolide, flurandrenolon acetone, hydrocortisone, meprednisone, methylprednisolon, paramethasone, prednisolon, prednisol, triamcinolon and triamcinolon acetone; narcotic agonists and antagonists such as apomorphine, buprenorphine, butorphanol, codein, cyclazocin, hydromorphon, ketobemidon, levallorphan, levorphanol, metazocin, morphine, nalbuphin, nalmefen, naloxon, nalorphine, naltrexon, oxycodon, oxymorphon and pentazocin; stimulants such as mazindol and pseudoephedrine; anaesthetics such as hydroxydion and propofol; β -receptor blockers such as acebutolol, albuterol, alprenolol, atenolol, betazolol, bucindolol, cartelolol, celiprolol, cetamolol, labetalol, levobunelol, metoprolol, metipranolol, nadolol, oxyprenolol, pindolol, propranolol and timolol; α -sympathomimetics such as adrenalin,

10001-334255



metaraminol, midodrin, norfenefrin, octapamine, oxedrin, oxilofrin, oximetazolin and phenylefrin; β -sympathomimetics such as bamethan, clenbuterol, fenoterol, hexoprenalin, isoprenalin, isoxsuprin, orciprenalin, reproterol, salbutamol and terbutalin; bronchodilators such as carbutoleol, dyphillin, etophyllin, fenoterol, pirbuterol, rimiterol and terbutalin; 5 cardiotonics such as digitoxin, dobutamin, etilefrin and prenalterol; antimycotics such as amphotericin B, chlorphenesin, nystatin and perimycin; anticoagulants such as acenocoumarol, dicoumarol, phenprocoumon and warfarin; vasodilators such as bamethan, dipyrimadol, diprophyllin, 10 isoxsuprin, vincamin and xantinol nicotinate; antihypocholesteremics such as compactin, eptastatin, mevinolin and simvastatin; miscellaneous drugs such as bromperidol (antipsychotic), dithranol (psoriasis) ergotamine (migraine) ivermectin (antihelminthic), metronidazole and secnizadole (antiprotozoals), nandrolon (anabolic), propafenon and quinadine (antiarythmics), quetiapine 15 (CNS), serotonin (neurotransmitter) and silybin (hepatic disturbance).

It is understood that the drug is not a GABA analog; L-Dopa, an L-aromatic amino acid decarboxylase inhibitor, a catechol O-methyl transferase inhibitor or derivatives thereof; a naturally occurring α -amino acid or an ester or 20 carboxamide of a naturally occurring α -amino acid; a polypeptide derived from a linear oligopeptide containing at least 3 α -amino acids; an oligonucleotide; a cyclophane derivative, a diethylenetriaminopentaacetate derivative, or paramagnetic ion chelates thereof; histamine or tyramine; 5-de-O-methylsporadicin; a bis-(2-chloroethyl)amine containing nitrogen mustard; an 25 HMG-CoA reductase inhibitor; a proline hydroxylase inhibitor; fluvalinate; or a steroid containing the carbon substructures of the following formulae:

106001-394266



“GABA analog” refers to a compound of one of the following formulae:



wherein:

R^4 is hydrogen, or R^4 and R^9 together with the atoms to which they are attached form a heterocyclic ring;

- 5 R^5 and R^6 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

R^7 and R^8 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, alkynyl, aryl, substituted aryl,

heteroaryl and substituted heteroaryl, or R^7 and R^8 together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclic or substituted heterocyclic ring;

5 R^9 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, alkynyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

R^{10} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, alkynyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

10 R^{11} is selected from the group consisting of carboxylic acid, carboxylic amide, carboxylic ester, sulfonamide, phosphonic acid, acidic heterocycle, sulfonic acid and hydroxamic acid;

$R^{4'}$ is hydrogen, or $R^{4'}$ and $R^{9'}$ together with the atoms to which they are attached form a heterocyclic ring;

15 $R^{5'}$ and $R^{6'}$ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

$R^{7'}$ and $R^{8'}$ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, alkynyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl, or $R^{7'}$ and $R^{8'}$ together with the atoms to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocyclic or substituted heterocyclic ring;

20 $R^{9'}$ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, alkynyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

25 $R^{10'}$ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, alkynyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl;

106001-39242660

R^{11'} is selected from the group consisting of carboxylic acid, carboxylic amide, carboxylic ester, sulfonamide, phosphonic acid, acidic heterocycle, sulfonic acid and hydroxamic acid.

“Catechol O-methyl transferase inhibitor” preferably refers to catechol
5 O-methyl transferase inhibitors such as entacapone, nitecapone and tolcapone optionally with one or two hydrogen atoms of two hydroxyl groups of the catechol group replaced with -C(O)R⁴⁴, -C(O)OR⁴⁵ and/or
-OCR⁴³R⁴⁴OC(O)R⁴⁵, wherein R⁴³ and R⁴⁴ independently are members selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl,
10 substituted cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl, or R⁴³ and R⁴⁴ together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocycle or substituted heterocyclic ring, R⁴⁵ is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl,
15 substituted aryl, aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl, or the OH group of the carboxyl moiety is replaced by -OR⁴⁴.

“An inhibitor of L-aromatic amino acid decarboxylase” preferably refers to L-aromatic amino acid decarboxylase inhibitors such as carbidopa and benzserazide optionally with a hydrogen atom of the amino or the hydrazido
20 group of the L-aromatic amino acid decarboxylase inhibitor replaced with -C(O)R¹⁰⁴, -C(O)OR¹⁰⁵ or an amino acid group, wherein R¹⁰⁴ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl, and R¹⁰⁵ is selected from the group consisting of alkyl,
25 substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl; and/or

optionally with one or two hydrogen atoms of the two –OH groups of the catechol or the three –OH groups of the pyrogallol group of the L-aromatic amino acid decarboxylase inhibitor are replaced with $-C(O)R^{104}$, $-C(O)OR^{105}$ and/or $-OCR^{103}R^{104}OC(O)R^{105}$ wherein R^{105} is defined as above, R^{103} and R^{104} independently are members selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl, or R^{103} and R^{104} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocycle or substituted heterocyclic ring; or optionally with two adjacent –OH groups of the catechol or pyrogallol group protected with a 5-membered cyclic carbonate or 2,3-dioxo-1,4-dioxane ortho fused with a benzene ring of the catechol or pyrogallol group; and/or the OH group of the carboxyl moiety is replaced by $-OR^{104}$.


“Derivatives of L-DOPA” preferably refers to L-DOPA molecules wherein:

a) a hydrogen atom of the amino group of the L-DOPA molecule is replaced with $-C(O)R^{114}$, $-C(O)OR^{115}$ or an amino acid group, wherein R^{114} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl, and R^{115} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl; and/or

b) one or two hydrogen atoms of the two –OH groups of the catechol group of the L-DOPA molecule are replaced with $-C(O)R^{114}$, $-C(O)OR^{115}$ and/or $-OCR^{113}R^{114}OC(O)R^{115}$ wherein R^{115} is defined as above, R^{113} and R^{114} independently are members selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl,


- aralkyl, substituted aralkyl, heteroaryl and substituted heteroaryl, or R^{113} and R^{114} together with the carbon atom to which they are attached form a cycloalkyl, substituted cycloalkyl, heterocycle or substituted heterocyclic ring, or the two -OH groups of the catechol group of the L-DOPA molecule are protected with a
- 5 5-membered cyclic carbonate or 2,3-dioxo-1,4-dioxane ortho fused with a benzene ring of the catechol group of the L-DOPA molecule; and/or
- c) the OH group of the carboxyl moiety is replaced by $-OR^{114}$.

“Acidic heterocycle” refers to a reprotonatable heterocycle having a pKa less than 7.0. Examples of such heterocycles include the following:



- 10 “Linear oligopeptide” refers to an amide oligomer comprising either a terminal amino group or a terminal carboxylic acid group or (preferably) both a terminal amino group and a terminal carboxylic acid group, which oligomer is formed by condensation of the terminal amino residue of at least one amino acid with the terminal carboxylic acid residue of at least a second amino acid. The
- 15 amino acids comprising the oligopeptide are either α -amino acids, β -amino acids, or a mixture of α -amino acids and β -amino acids. Note that when an α -amino acid additionally contains either a β -amino group or a β -carboxylic acid group (e.g. as in aspartic acid) a linear oligopeptide formed from such an amino acid is intended to imply that it is the α -amine or α -carboxylic acid moiety (or
- 20 both) of such residue that is involved in amide formation.

“Derived from a compound” or “derivative” refers to a moiety that is structurally related to such a compound. The structure of the moiety is identical to the compound except at 1 or 2 positions. At these positions either a hydrogen atom attached to a heteroatom, or a hydroxyl moiety of a carboxylic, phosphonic, phosphoric or sulfonic acid group has been replaced with a covalent bond that serves as a point of attachment to another moiety. For example, the moiety:



is derived from a linear oligopeptide comprising glycine and the drug gabapentin. In this moiety, a hydrogen atom has been replaced with a covalent bond. “Derived from a linear oligopeptide” is meant to specifically denote that the point of attachment is either the terminal amino group or the terminal acid group of the oligopeptide.

The term “systemic bioavailability” refers to the rate and extent of systemic exposure to a drug or a metabolite thereof as reflected by the area under the systemic blood concentration versus time curve.

“Translocation across the intestinal wall” refers to movement of a drug or drug conjugate by a passive or active mechanism, or both, across an epithelial cell membrane of any region of the gastrointestinal tract.

“Active metabolite of a drug” refers to products of *in vivo* modification of the drug which have therapeutic or prophylactic effect.

“Therapeutic or prophylactic blood concentrations” refers to systemic exposure to a sufficient concentration of a drug or an active metabolite thereof over a sufficient period of time to effect disease therapy or to prevent the onset or reduce the severity of a disease in the treated animal.

5 “Sustained release” refers to release of a drug or an active metabolite thereof into the systemic circulation over a prolonged period of time relative to that achieved by oral administration of a conventional formulation of the drug.

10 “Tissue of the enterohepatic circulation” refers to the blood, plasma, intestinal contents, intestinal cells, liver cells, biliary tract or any fraction, suspension, homogenate, extract or preparation thereof.

“Conjugating” refers to the formation of a covalent bond.

“Bile acid transport system” refers to any membrane transporter protein capable of causing a bile acid or a derivative thereof to be translocated across a membrane of a cell of the gastrointestinal tract or liver.

15 “Active transport or active transport mechanism” refers to the movement of molecules across cellular membranes that: a) is directly or indirectly dependent on an energy mediated process (i.e. driven by ATP hydrolysis, ion gradient, etc); or b) occurs by facilitated diffusion mediated by interaction with specific transporter proteins; or c) occurs through a modulated solute channel.

20 “A moiety selected to permit a compound of formula (I) to be translocated across the intestinal wall of an animal via the bile acid transport system” refers to compounds which, when conjugated to the drug (directly or via a linker moiety), are translocated across the intestinal wall via the bile acid

105001-39242660

transport system. Evaluation of which candidate compounds can be so translocated across the intestinal wall can be conducted by the *in vitro* assay set forth in Example 3 below.

5 “Practical dosage regimen” refers to a schedule of drug administration that is practical for a patient to comply with. For human patients, a practical dosage regimen for an orally administered drug is likely to be an aggregate dose of less than 10 g/day.

10 “Amino-protecting group” or “amino-blocking group” refers to any group which when bound to one or more amino groups prevents reactions from occurring at these amino groups and which protecting groups can be removed by conventional chemical steps to reestablish the amino group. The particular removable blocking group is not critical and preferred amino blocking groups include, by way of example only, t-butyloxycarbonyl (t-BOC),
15 benzyloxycarbonyl (CBZ), and the like.

20 “Carboxyl-protecting group” or “carboxyl-blocking group” refers to any group which when bound to one or more carboxyl groups prevents reactions from occurring at these groups and which protecting groups can be removed by conventional chemical steps to reestablish the carboxyl group. The particular removable blocking group is not critical and preferred carboxyl blocking groups include, by way of example only, esters of the formula $-COOR$ where R” is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, alkaryl, substituted alkaryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and
25 substituted heterocyclic.

00445-1009
T06001-894450

"Alkyl" refers to alkyl groups preferably having from 1 to 20 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, t-butyl, n-heptyl, octyl, dodecyl and the like.

5 "Substituted alkyl" refers to an alkyl group, preferably of from 1 to 20 carbon atoms, having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxylaryl, substituted aryloxyaryl, cyano, halogen, 10 hydroxyl, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted 15 thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted aryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, 20 oxythiocarbonylamino, $-\text{OS}(\text{O})_2\text{-alkyl}$, $-\text{OS}(\text{O})_2\text{-substituted alkyl}$, $-\text{OS}(\text{O})_2\text{-aryl}$, $-\text{OS}(\text{O})_2\text{-substituted aryl}$, $-\text{OS}(\text{O})_2\text{-heteroaryl}$, $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$, $-\text{OS}(\text{O})_2\text{-heterocyclic}$, $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$, $-\text{OSO}_2\text{-NRR}$ where R is hydrogen or alkyl, $-\text{NRS}(\text{O})_2\text{-alkyl}$, $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$, $-\text{NRS}(\text{O})_2\text{-aryl}$, $-\text{NRS}(\text{O})_2\text{-substituted aryl}$, $-\text{NRS}(\text{O})_2\text{-heteroaryl}$, $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$, $-\text{NRS}(\text{O})_2\text{-heterocyclic}$, $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$, $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$, $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$, $-\text{NRS}(\text{O})_2\text{-NR-aryl}$, $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$, $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$, $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$, $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$, $-\text{NRS}(\text{O})_2\text{-NR-substituted}$

000001-0924660

heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkyl/substituted alkyl groups substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

"Alkoxy" refers to the group "alkyl-O-" which includes, by way of example, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

"Substituted alkoxy" refers to the group "substituted alkyl-O-".

"Acyl" refers to the groups H-C(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, alkenyl-C(O)-, substituted alkenyl-C(O)-, alkynyl-C(O)-, substituted alkynyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, aryl-C(O)-, substituted aryl-C(O)-, heteroaryl-C(O)-, substituted heteroaryl-C(O), heterocyclic-C(O)-, and substituted heterocyclic-C(O)- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Acylamino" refers to the group -C(O)NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Thiocarbonylamino" refers to the group -C(S)NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and where each R is joined to form, together with the nitrogen atom a heterocyclic or substituted heterocyclic ring wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Acyloxy" refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-, alkenyl-C(O)O-, substituted alkenyl-C(O)O-, alkynyl-C(O)O-, substituted alkynyl-C(O)O-, aryl-C(O)O-, substituted aryl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, heteroaryl-C(O)O-, substituted heteroaryl-C(O)O-, heterocyclic-C(O)O-, and substituted heterocyclic-C(O)O- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl,

cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Alkenyl" refers to alkenyl group preferably having from 2 to 20 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkenyl unsaturation.

"Substituted alkenyl" refers to alkenyl groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, $-\text{OS}(\text{O})_2\text{-alkyl}$, $-\text{OS}(\text{O})_2\text{-substituted alkyl}$, $-\text{OS}(\text{O})_2\text{-aryl}$, $-\text{OS}(\text{O})_2\text{-substituted aryl}$, $-\text{OS}(\text{O})_2\text{-heteroaryl}$, $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$, $-\text{OS}(\text{O})_2\text{-heterocyclic}$, $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$, $-\text{OSO}_2\text{-NRR}$ where R is hydrogen or alkyl, $-\text{NRS}(\text{O})_2\text{-alkyl}$, $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$, $-\text{NRS}(\text{O})_2\text{-aryl}$, $-\text{NRS}(\text{O})_2\text{-substituted aryl}$,

-NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -
NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl,
-NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl,
-NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-
5 substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted
heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and
di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted
arylamino, mono- and di-heteroarylamino, mono- and di-substituted
heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted
10 heterocyclic amino, unsymmetric di-substituted amines having different
substituents selected from the group consisting of alkyl, substituted alkyl, aryl,
substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted
heterocyclic and substituted alkenyl groups having amino groups blocked by
conventional blocking groups such as Boc, Cbz, formyl, and the like or
15 alkenyl/substituted alkenyl groups substituted with -SO₂-alkyl, -SO₂-substituted
alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl,
-SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl,
-SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic
and -SO₂NRR where R is hydrogen or alkyl.

20 "Alkenyloxy" refers to the group -O-alkenyl.

"Substituted alkenyloxy" refers to the group -O-substituted alkenyloxy.

"Alkynyl" refers to alkynyl group preferably having from 2 to 20 carbon
atoms and more preferably 3 to 6 carbon atoms and having at least 1 and
25 preferably from 1-2 sites of alkynyl unsaturation.

"Substituted alkynyl" refers to alkynyl groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted

heteroaryl-amino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

"Alkylene" refers to a divalent alkylene group preferably having from 1 to 20 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methylene (-CH₂-), ethylene (-CH₂CH₂-), the propylene isomers (e.g., -CH₂CH₂CH₂- and -CH(CH₃)CH₂-) and the like.

"Substituted alkylene" refers to alkylene groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted

thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl,
substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy,
substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,
heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino,
5 oxythiocarbonylamino, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -
OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -
OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is
hydrogen or alkyl, -NRS(O)₂-alkyl,
- NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl,
10 -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -
NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl,
- NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl,
- NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-
substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted
15 heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and
di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted
arylamino, mono- and di-heteroarylamino, mono- and di-substituted
heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted
heterocyclic amino, unsymmetric di-substituted amines having different
20 substituents selected from the group consisting of alkyl, substituted alkyl, aryl,
substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted
heterocyclic and substituted alkenyl groups having amino groups blocked by
conventional blocking groups such as Boc, Cbz, formyl, and the like or
alkenyl/substituted alkenyl groups substituted with -SO₂-alkyl, -SO₂-substituted
25 alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl,
-SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -
SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and
-SO₂NRR where R is hydrogen or alkyl.

TOP SECRET 6924660

"Alkenylene" refers to a divalent alkenylene group preferably having from 2 to 20 carbon atoms and more preferably 1 to 6 carbon atoms and having from 1 to 2 sites of alkenyl unsaturation. This term is exemplified by groups such as ethenylene ($-\text{CH}=\text{CH}-$), propenylene ($-\text{CH}_2\text{CH}=\text{CH}-$), and the like.

- 5 "Substituted alkenylene" refers to alkenylene groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, 10 aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, 15 guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, 20 heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, $-\text{OS}(\text{O})_2$ -alkyl, $-\text{OS}(\text{O})_2$ -substituted alkyl, $-\text{OS}(\text{O})_2$ -aryl, $-\text{OS}(\text{O})_2$ -substituted aryl, $-\text{OS}(\text{O})_2$ -heteroaryl, $-\text{OS}(\text{O})_2$ -substituted heteroaryl, $-\text{OS}(\text{O})_2$ -heterocyclic, $-\text{OS}(\text{O})_2$ -substituted heterocyclic, $-\text{OSO}_2$ -NRR where R is hydrogen or alkyl, $-\text{NRS}(\text{O})_2$ -alkyl, 25 $-\text{NRS}(\text{O})_2$ -substituted alkyl, $-\text{NRS}(\text{O})_2$ -aryl, $-\text{NRS}(\text{O})_2$ -substituted aryl, $-\text{NRS}(\text{O})_2$ -heteroaryl, $-\text{NRS}(\text{O})_2$ -substituted heteroaryl, $-\text{NRS}(\text{O})_2$ -heterocyclic, $-\text{NRS}(\text{O})_2$ -substituted heterocyclic, $-\text{NRS}(\text{O})_2$ -NR-alkyl, $-\text{NRS}(\text{O})_2$ -NR-substituted alkyl, $-\text{NRS}(\text{O})_2$ -NR-aryl,

-NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and substituted alkenyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkenyl/substituted alkenyl groups substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

"Alkynylene" refers to a divalent alkynylene group preferably having from 2 to 20 carbon atoms and more preferably 1 to 6 carbon atoms and having from 1 to 2 sites of alkynyl unsaturation. This term is exemplified by groups such as ethynylene, propynylene and the like.

"Substituted alkynylene" refers to alkynylene groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-

substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,
 carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,
 carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino,
 guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted
 5 thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted
 thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl,
 substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy,
 substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,
 heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino,
 10 oxythiocarbonylamino, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -
 OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -
 OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is
 hydrogen or alkyl, -NRS(O)₂-alkyl,
 -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl,
 15 -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -
 NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl,
 -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted
 aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-
 NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or
 20 alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono-
 and di-arylamino, mono- and di-substituted arylamino, mono- and di-
 heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-
 heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric
 di-substituted amines having different substituents selected from the group
 25 consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,
 substituted heteroaryl, heterocyclic, substituted heterocyclic and substituted
 alkenyl groups having amino groups blocked by conventional blocking groups
 such as Boc, Cbz, formyl, and the like or alkenyl/substituted alkenyl groups
 substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-

T0600T-89242660

substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

5 "Amidino" refers to the group H₂NC(=NH)- and the term "alkylamidino" refers to compounds having 1 to 3 alkyl groups (e.g., alkylHNC(=NH)-).

"Thioamidino" refers to the group RSC(=NH)- where R is hydrogen or alkyl.

10 "Aminoacyl" refers to the groups -NRC(O)alkyl, -NRC(O)substituted alkyl, -NRC(O)cycloalkyl, -NRC(O)substituted cycloalkyl, -NRC(O)alkenyl, -NRC(O)substituted alkenyl, -NRC(O)alkynyl, -NRC(O)substituted alkynyl, -NRC(O)aryl, -NRC(O)substituted aryl, -NRC(O)heteroaryl, -NRC(O)substituted heteroaryl, -NRC(O)heterocyclic, and
15 -NRC(O)substituted heterocyclic where R is hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminocarbonyloxy" refers to the groups -NRC(O)O-alkyl, -NRC(O)O-substituted alkyl, -NRC(O)O-alkenyl, -NRC(O)O-substituted alkenyl, -NRC(O)O-alkynyl, -NRC(O)O-substituted alkynyl, -NRC(O)O-cycloalkyl, -NRC(O)O-substituted cycloalkyl, -NRC(O)O-aryl, -NRC(O)O-substituted aryl, -NRC(O)O-heteroaryl, -NRC(O)O-substituted heteroaryl, -NRC(O)O-heterocyclic, and -NRC(O)O-substituted heterocyclic where R is
20 hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted
25

0949260

alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Oxycarbonylamino" refers to the groups -OC(O)NH₂, -OC(O)NRR, -OC(O)NR-alkyl, -OC(O)NR-substituted alkyl, -OC(O)NR-alkenyl, -OC(O)NR-substituted alkenyl, -OC(O)NR-alkynyl, -OC(O)NR-substituted alkynyl, -OC(O)NR-cycloalkyl, -OC(O)NR-substituted cycloalkyl, -OC(O)NR-aryl, -OC(O)NR-substituted aryl, -OC(O)NR-heteroaryl, -OC(O)NR-substituted heteroaryl, -OC(O)NR-heterocyclic, and -OC(O)NR-substituted heterocyclic where R is hydrogen, alkyl or where each R is joined to form, together with the nitrogen atom a heterocyclic or substituted heterocyclic ring and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Oxythiocarbonylamino" refers to the groups -OC(S)NH₂, -OC(S)NRR, -OC(S)NR-alkyl, -OC(S)NR-substituted alkyl, -OC(S)NR-alkenyl, -OC(S)NR-substituted alkenyl, -OC(S)NR-alkynyl, -OC(S)NR-substituted alkynyl, -OC(S)NR-cycloalkyl, -OC(S)NR-substituted cycloalkyl, -OC(S)NR-aryl, -OC(S)NR-substituted aryl, -OC(S)NR-heteroaryl, -OC(S)NR-substituted heteroaryl, -OC(S)NR-heterocyclic, and -OC(S)NR-substituted heterocyclic where R is hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminocarbonylamino" refers to the groups -NRC(O)NRR, -NRC(O)NR-alkyl, -NRC(O)NR-substituted alkyl, -NRC(O)NR-alkenyl, -NRC(O)NR-substituted alkenyl, -NRC(O)NR-alkynyl, -NRC(O)NR-substituted alkynyl, -NRC(O)NR-aryl, -NRC(O)NR-substituted aryl, -NRC(O)NR-cycloalkyl, -NRC(O)NR-substituted cycloalkyl, -NRC(O)NR-heteroaryl, and -NRC(O)NR-substituted heteroaryl, -NRC(O)NR-heterocyclic, and -NRC(O)NR-substituted heterocyclic where each R is independently hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminothiocabonylamino" refers to the groups -NRC(S)NRR, -NRC(S)NR-alkyl, -NRC(S)NR-substituted alkyl, -NRC(S)NR-alkenyl, -NRC(S)NR-substituted alkenyl, -NRC(S)NR-alkynyl, -NRC(S)NR-substituted alkynyl, -NRC(S)NR-aryl, -NRC(S)NR-substituted aryl, -NRC(S)NR-cycloalkyl, -NRC(S)NR-substituted cycloalkyl, -NRC(S)NR-heteroaryl, and -NRC(S)NR-substituted heteroaryl, -NRC(S)NR-heterocyclic, and -NRC(S)NR-substituted heterocyclic where each R is independently hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aryl" or "Ar" refers to a monovalent unsaturated aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl or anthryl) which condensed rings may or may not be aromatic (e.g., 2-benzoxazolinone, 2H-1,4-benzoxazin-3(4H)-one-7yl, and the like). Preferred aryls include phenyl and naphthyl.

Substituted aryl refers to aryl groups which are substituted with from 1 to 3 substituents selected from the group consisting of hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy, aminocarbonylamino, aminothiocarbonylamino, aryl, substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -S(O)₂-alkyl, -S(O)₂-substituted alkyl, -S(O)₂-cycloalkyl, -S(O)₂-substituted cycloalkyl, -S(O)₂-alkenyl, -S(O)₂-substituted alkenyl, -S(O)₂-aryl, -S(O)₂-substituted aryl, -S(O)₂-heteroaryl, -S(O)₂-substituted heteroaryl, -S(O)₂-heterocyclic, -S(O)₂-substituted heterocyclic, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted

heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -
 NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -
 NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -
 NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-
 5 substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-
 NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-
 heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or
 alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono-
 and di-arylamino, mono- and di-substituted arylamino, mono- and di-
 10 heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-
 heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric
 di-substituted amines having different substituents selected from the group
 consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,
 substituted heteroaryl, heterocyclic and substituted heterocyclic and amino
 15 groups on the substituted aryl blocked by conventional blocking groups such as
 Boc, Cbz, formyl, and the like or substituted with -SO₂NRR where R is
 hydrogen or alkyl.

"Arylene" refers to a divalent unsaturated aromatic carbocyclic group of
 from 6 to 14 carbon atoms having a single ring (e.g., phenylene) or multiple
 20 condensed rings (e.g., naphthylene or anthrylene) which condensed rings may or
 may not be aromatic. Preferred arylenes include phenylene and naphthylene.
 Substituted arylene refers to arylene groups which are substituted with from 1 to
 3 substituents selected from the group consisting of hydroxy, acyl, acylamino,
 thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy, substituted alkoxy,
 25 alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, amidino, alkylamidino,
 thioamidino, amino, aminoacyl, aminocarbonyloxy, aminocarbonylamino,
 aminothiocarbonylamino, aryl, substituted aryl, aryloxy, substituted aryloxy,
 cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,

0947681000T" 89242660

heterocyclyloxy, substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -S(O)₂-alkyl, -S(O)₂-substituted alkyl, -S(O)₂-cycloalkyl, -S(O)₂-substituted cycloalkyl, -S(O)₂-alkenyl, -S(O)₂-substituted alkenyl, -S(O)₂-aryl, -S(O)₂-substituted aryl, -S(O)₂-heteroaryl, -S(O)₂-substituted heteroaryl, -S(O)₂-heterocyclic, -S(O)₂-substituted heterocyclic, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group

094450 106001-394650

consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,
substituted heteroaryl, heterocyclic and substituted heterocyclic and amino
groups on the substituted aryl blocked by conventional blocking groups such as
Boc, Cbz, formyl, and the like or substituted with -SO₂NRR where R is
5 hydrogen or alkyl.

"Aryloxy" refers to the group aryl-O- which includes, by way of
example, phenoxy, naphthoxy, and the like.

"Substituted aryloxy" refers to substituted aryl-O- groups.

"Aryloxyaryl" refers to the group -aryl-O-aryl.

10 "Substituted aryloxyaryl" refers to aryloxyaryl groups substituted with
from 1 to 3 substituents on either or both aryl rings selected from the group
consisting of hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl,
substituted alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl,
alkynyl, substituted alkynyl, amidino, alkylamidino, thioamidino, amino,
15 aminoacyl, aminocarbonyloxy, aminocarbonylamino, aminothiocarbonylamino,
aryl, substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted
cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy,
substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl,
carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-
20 substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl,
carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano,
thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,
thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted
thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl,
25 substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl,

106001-2924660

substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy,
substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,
heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino,
oxythiocarbonylamino, -S(O)₂-alkyl, -S(O)₂-substituted alkyl, -S(O)₂-cycloalkyl,
5 -S(O)₂-substituted cycloalkyl, -S(O)₂-alkenyl, -S(O)₂-substituted alkenyl, -S(O)₂-
aryl, -S(O)₂-substituted aryl, -S(O)₂-heteroaryl, -S(O)₂-substituted heteroaryl, -
S(O)₂-heterocyclic, -S(O)₂-substituted heterocyclic, -OS(O)₂-alkyl, -OS(O)₂-
substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl,
-OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted
10 heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl,
-NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl,
-NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -
NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-
substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-
15 NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-
heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or
alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono-
and di-arylamino, mono- and di-substituted arylamino, mono- and di-
heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-
20 heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric
di-substituted amines having different substituents selected from the group
consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,
substituted heteroaryl, heterocyclic and substituted heterocyclic and amino
groups on the substituted aryl blocked by conventional blocking groups such as
25 Boc, Cbz, formyl, and the like or substituted with -SO₂NRR where R is
hydrogen or alkyl.

"Cycloalkyl" refers to cyclic alkyl groups of from 3 to 8 carbon atoms
having a single cyclic ring including, by way of example, cyclopropyl,

cyclobutyl, cyclopentyl, cyclooctyl and the like. Excluded from this definition are multi-ring alkyl groups such as adamantanyl, etc.

"Cycloalkyloxy" refers to -O-cycloalkyl.

5 "Cycloalkenyl" refers to cyclic alkenyl groups of from 3 to 8 carbon atoms having a single cyclic ring.

"Cycloalkenyloxy" refers to -O-cycloalkenyl.

10 "Substituted cycloalkyl" and "substituted cycloalkenyl" refers to an cycloalkyl or cycloalkenyl group, preferably of from 3 to 8 carbon atoms, having from 1 to 5 substituents selected from the group consisting of oxo (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-
15 cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted
20 thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -
25 OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -

OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -

5 NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted

10 arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted

15 heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted

20 heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

"Substituted cycloalkyloxy" and "substituted cycloalkenyloxy" refers to -O-substituted cycloalkyl and -O-substituted cycloalkenyloxy respectively.

"Cycloalkylene" refers to divalent cyclic alkylene groups of from 3 to 8

25 carbon atoms having a single cyclic ring including, by way of example, cyclopropylene, cyclobutylene, cyclopentylene, cyclooctylene and the like.

"Cycloalkenylene" refers to a divalent cyclic alkenylene groups of from 3 to 8 carbon atoms having a single cyclic ring.

"Substituted cycloalkylene" and "substituted cycloalkenylene" refers to a cycloalkylene or cycloalkenylene group, preferably of from 3 to 8 carbon atoms, having from 1 to 5 substituents selected from the group consisting of oxo (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-

substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

"Cycloalkoxy" refers to -O-cycloalkyl groups.

"Substituted cycloalkoxy" refers to -O-substituted cycloalkyl groups.

"Guanidino" refers to the groups -NRC(=NR)NRR, -NRC(=NR)NR-alkyl, -NRC(=NR)NR-substituted alkyl, -NRC(=NR)NR-alkenyl, -NRC(=NR)NR-substituted alkenyl, -NRC(=NR)NR-alkynyl, -NRC(=NR)NR-substituted alkynyl, -NRC(=NR)NR-aryl, -NRC(=NR)NR-substituted aryl, -NRC(=NR)NR-cycloalkyl, -NRC(=NR)NR-heteroaryl, -NRC(=NR)NR-substituted heteroaryl, -NRC(=NR)NR-heterocyclic, and -NRC(=NR)NR-substituted heterocyclic where each R is independently hydrogen and alkyl as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted

alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"*N,N*-Dimethylcarbamyloxy" refers to the group $-\text{OC}(\text{O})\text{N}(\text{CH}_3)_2$.

5 "Guanidinosulfone" refers to the groups $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -alkyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -substituted alkyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -alkenyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -substituted alkenyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -alkynyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -substituted alkynyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -aryl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -substituted aryl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -cycloalkyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -substituted cycloalkyl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -heteroaryl, and $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -substituted heteroaryl, $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -heterocyclic, and $-\text{NRC}(=\text{NR})\text{NRSO}_2$ -substituted heterocyclic where each R is independently hydrogen and alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

10

15

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo and preferably is either chloro or bromo.

"Heteroaryl" refers to an aromatic carbocyclic group of from 2 to 10 carbon atoms and 1 to 4 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur within the ring. Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indoliziny or benzothienyl). Preferred heteroaryls include pyridyl, pyrrolyl, indolyl and furyl.

20

"Substituted heteroaryl" refers to heteroaryl groups which are substituted with from 1 to 3 substituents selected from the group consisting of hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy, aminocarbonylamino, aminothiocarbonylamino, aryl, substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -S(O)₂-alkyl, -S(O)₂-substituted alkyl, -S(O)₂-cycloalkyl, -S(O)₂-substituted cycloalkyl, -S(O)₂-alkenyl, -S(O)₂-substituted alkenyl, -S(O)₂-aryl, -S(O)₂-substituted aryl, -S(O)₂-heteroaryl, -S(O)₂-substituted heteroaryl, -S(O)₂-heterocyclic, -S(O)₂-substituted heterocyclic, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -

106001-89242660

106907-094760

NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and amino groups on the substituted aryl blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or substituted with -SO₂NRR where R is hydrogen or alkyl.

"Heteroarylene" refers to a divalent aromatic carbocyclic group of from 2 to 10 carbon atoms and 1 to 4 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur within the ring. Such heteroarylene groups can have a single ring (e.g., pyridylene or furylene) or multiple condensed rings (e.g., indolizinylene or benzothienylene). Preferred heteroarylenes include pyridylene, pyrrolylene, indolyene and furylene.

"Substituted heteroarylene" refers to heteroarylene groups which are substituted with from 1 to 3 substituents selected from the group consisting of hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy, aminocarbonylamino, aminothiocabonylamino, aryl, substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl,

carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,

5 thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,

10 heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -S(O)₂-alkyl, -S(O)₂-substituted alkyl, -S(O)₂-cycloalkyl, -S(O)₂-substituted cycloalkyl, -S(O)₂-alkenyl, -S(O)₂-substituted alkenyl, -S(O)₂-aryl, -S(O)₂-substituted aryl, -S(O)₂-heteroaryl, -S(O)₂-substituted heteroaryl, -S(O)₂-heterocyclic, -S(O)₂-substituted heterocyclic, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,

25

1066007-89242660

substituted heteroaryl, heterocyclic and substituted heterocyclic and amino groups on the substituted aryl blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or substituted with -SO₂NRR where R is hydrogen or alkyl.

- 5 "Heteroaryloxy" refers to the group -O-heteroaryl and "substituted heteroaryloxy" refers to the group -O-substituted heteroaryl.

- 10 "Heterocycle" or "heterocyclic" refers to a saturated or unsaturated group having a single ring or multiple condensed rings, from 1 to 10 carbon atoms and from 1 to 4 hetero atoms selected from the group consisting of nitrogen, sulfur or oxygen within the ring wherein, in fused ring systems, one or more the rings can be aryl or heteroaryl.

- 15 "Substituted heterocyclic" refers to heterocycle groups which are substituted with from 1 to 3 substituents selected from the group consisting of oxo (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, 20 carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, 25 substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy,

substituted heteroaryloxy, -C(O)O-aryl, -C(O)O-substituted aryl, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

Examples of heterocycles and heteroaryls include, but are not limited to, azetidine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, dihydroindole, indazole, purine,

quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, phthalimide, 5 1,2,3,4-tetrahydroisoquinoline, 4,5,6,7-tetrahydrobenzo[b]thiophene, thiazole, thiazolidine, thiophene, benzo[b]thiophene, morpholinyl, thiomorpholinyl (also referred to as thiamorpholinyl), piperidinyl, pyrrolidine, tetrahydrofuranyl, and the like.

"Heterocyclene" refers to a divalent saturated or unsaturated group 10 having a single ring or multiple condensed rings, from 1 to 10 carbon atoms and from 1 to 4 hetero atoms selected from the group consisting of nitrogen, sulfur or oxygen within the ring wherein, in fused ring systems, one or more the rings can be aryl or heteroaryl.

"Substituted heterocyclene" refers to heterocyclene groups which are 15 substituted with from 1 to 3 substituents selected from the group consisting of oxo (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted 20 aryloxyaryl, halogen, hydroxyl, cyano, nitro, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted 25 thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic,

09974768-100001

substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, -C(O)O-aryl, -C(O)O-substituted aryl, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)₂-alkyl, -OS(O)₂-substituted alkyl, -OS(O)₂-aryl, -OS(O)₂-substituted aryl, -OS(O)₂-heteroaryl, -OS(O)₂-substituted heteroaryl, -OS(O)₂-heterocyclic, -OS(O)₂-substituted heterocyclic, -OSO₂-NRR where R is hydrogen or alkyl, -NRS(O)₂-alkyl, -NRS(O)₂-substituted alkyl, -NRS(O)₂-aryl, -NRS(O)₂-substituted aryl, -NRS(O)₂-heteroaryl, -NRS(O)₂-substituted heteroaryl, -NRS(O)₂-heterocyclic, -NRS(O)₂-substituted heterocyclic, -NRS(O)₂-NR-alkyl, -NRS(O)₂-NR-substituted alkyl, -NRS(O)₂-NR-aryl, -NRS(O)₂-NR-substituted aryl, -NRS(O)₂-NR-heteroaryl, -NRS(O)₂-NR-substituted heteroaryl, -NRS(O)₂-NR-heterocyclic, -NRS(O)₂-NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-alkenyl, -SO₂-substituted alkenyl, -SO₂-cycloalkyl, -SO₂-substituted cycloalkyl, -SO₂-aryl, -SO₂-substituted aryl, -SO₂-heteroaryl, -SO₂-substituted heteroaryl, -SO₂-heterocyclic, -SO₂-substituted heterocyclic and -SO₂NRR where R is hydrogen or alkyl.

"Heterocyclyloxy" refers to the group -O-heterocyclic and "substituted heterocyclyloxy" refers to the group -O-substituted heterocyclic.

"Thiol" refers to the group -SH.

"Thioalkyl" refers to the groups -S-alkyl

"Substituted thioalkyl" refers to the group -S-substituted alkyl.

"Thiocycloalkyl" refers to the groups -S-cycloalkyl.

5 "Substituted thiocycloalkyl" refers to the group -S-substituted cycloalkyl.

"Thioaryl" refers to the group -S-aryl and "substituted thioaryl" refers to the group -S-substituted aryl.

"Thioheteroaryl" refers to the group -S-heteroaryl and "substituted thioheteroaryl" refers to the group -S-substituted heteroaryl.

10 "Thioheterocyclic" refers to the group -S-heterocyclic and "substituted thioheterocyclic" refers to the group -S-substituted heterocyclic.

"Amino" refers to the -NH_2 group.

15 "Substituted amino" refers to the -NR'R'' group wherein R' and R'' are independently hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic or where R' and R'', together with the nitrogen atom pendent thereto, form a heterocyclic ring.

09974758-10994
106007-89242660

"Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts of a compound of Formulae (I), which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like.

Utility

The compounds and methods described herein provide for the sustained release of drugs or active metabolites thereof relative to dosing with the parent drug itself. In this regard, enterohepatic recycling of the bile acid conjugates creates a reservoir for the active agent.

General Synthetic Scheme

Compounds of this invention can be made by the methods depicted in the reaction schemes shown below.

The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, Wisconsin, USA), Bachem (Torrance, California, USA), Emka-Chemie, or Sigma (St. Louis, Missouri, USA) or are prepared by methods known to those skilled in the art following procedures set forth in references such as Fieser and Fieser's Reagents for Organic Synthesis, Volumes 1-15 (John Wiley and Sons, 1991); Rodd's Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989), Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991), March's Advanced Organic Chemistry, (John Wiley and Sons, 4th Edition), and Larock's Comprehensive Organic Transformations (VCH Publishers Inc., 1989).


These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure.

- 5 The starting materials and the intermediates of the reaction may be isolated and purified if desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography, and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.

10 Preparation of compounds of Formula (I)


Schemes A and B describe alternative methods to prepare the compounds of Formula (I).

- 15 Compounds of formula (I-1) where X, R¹ and R² are hydroxy, Z is a substituted alkyl group containing a group of the formula -M-Q^{x'}, M is -CH₂CH₂C(O)-, Q^{x'} is of the structure -A^{x'}-D', A^{x'} is derived from an α-amino acid, and D' is a drug of formula (b):



can be prepared as illustrated and described in Scheme A below.


Scheme A



In reference to Scheme A, commercially available cholic acid (A) is treated with an activating agent, such as ethylchloroformate in the presence of diisopropylethylamine, to provide an activated anhydride. The activated anhydride is reacted with amino acid (a) forming acid (B). Activation of (B) using an activating agent and condensation with drug (b) affords a compound of formula (I-1) as shown.


One of ordinary skill in the art will appreciate that cholic acid (A) could alternatively be reacted with amino acid (a) in the presence of a coupling agent (e.g., dicyclohexylcarbodiimide) to provide acid (B). Alternatively, cholic acid (A) could be coupled with the dipeptide (or depsipeptide) resulting from first coupling amino acid (a) to compound (b).

A compound of formula (I-2) where R^1 and R^2 are hydroxy, Z is $-\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, X is $\text{Q}^x-\text{G}-$, G is $-\text{O}-$, Q^x is $-\text{D}-\text{A}^x$, A^x is derived from an α -amino acid, and D is a drug of formula (b):



can be prepared as illustrated and described in Scheme B below.

Scheme B



In reference to Scheme B, compound (C), wherein R^{100} is a carboxyl protecting group, is coupled with activated acid derivative (c), wherein R^{101} is a suitable leaving group (*e.g.*, $-\text{OC}(\text{O})\text{OEt}$). Selective deprotection of the amino or hydroxyl protecting group (*i.e.*, Pg) provides compound (D). Activation of carboxylic acid (d) (*e.g.*, treatment with ethyl chloroformate in the presence of a

tertiary amine) followed by addition to (D) then removal of protecting groups yields the desired compound of formula (I-2).

One of ordinary skill in the art will appreciate that one could also use a coupling agent (e.g. a carbodiimide) *in situ* to form the amide or ester bonds above (*i.e.*, bond between (C) and (c) or between (D) and (d)) rather than condensing a preformed activated acid derivative (*i.e.*, (c) or (d)).

Analogous synthetic methods can be used in the preparation of compounds of formula (II) and (III).

Additionally, Figures 4-8 and Examples 1-8 below describe in detail the synthesis and biological activity of various bile acid-derived compounds.

Pharmaceutical Formulations

When employed as pharmaceuticals, the compounds of formula (I), (II) or (III) are usually administered in the form of pharmaceutical compositions. These compounds can be administered by a variety of routes including oral, rectal, subcutaneous, intravenous, intramuscular and intranasal. These compounds are effective as both injectable and oral compositions. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.

This invention also includes pharmaceutical compositions which contain, as the active ingredient, one or more of the compounds of formula (I) above associated with pharmaceutically acceptable carriers. In making the compositions of this invention, the active ingredient is usually mixed with an excipient, diluted by an excipient or enclosed within such a carrier which can be in the form of a capsule, sachet, paper or other container. When the excipient

5 serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

10 In preparing a formulation, it may be necessary to mill the active compound to provide the appropriate particle size prior to combining with other ingredients. If the active compound is substantially insoluble, it ordinarily is milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size is normally adjusted by milling to provide a substantially uniform distribution in the formulation, *e.g.* about 40 mesh.

15 Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, 20 magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxybenzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing 25 procedures known in the art.

5 The compositions are preferably formulated in a unit dosage form, each dosage containing from about 0.1 to about 5000 mg, more usually about 10 to about 2000 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other animals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

10 The active compound is effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It, will be understood, however, that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

15 For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, 0.1 mg to about 2 g of the active ingredient of the present invention.

25 The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged

action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable solvents may be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device may be attached to a face masks tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of this invention. Unless otherwise stated, all temperatures are in degrees Celsius.

EXAMPLES

5 In the examples below, the following abbreviations have the following meanings. If an abbreviation is not defined, it has its generally accepted meaning.

	Atm	=	atmosphere
	Boc	=	<i>tert</i> -butoxycarbonyl
10	Cbz	=	carbobenzyloxy
	CPM	=	counts per minute
	DIC	=	diisopropylcarbodiimide
	DMAP	=	4- <i>N,N</i> -dimethylaminopyridine
	DMEM	=	Dulbecco's minimum eagle medium
15	DMF	=	<i>N,N</i> -dimethylformamide
	DMSO	=	dimethylsulfoxide
	Fmoc	=	9-fluorenylmethyloxycarbonyl
	g	=	gram
	h	=	hour
20	HBSS	=	Hank's buffered saline solution
	IBAT	=	intestinal bile acid transporter
	L	=	liter
	LBAT	=	liver bile acid transporter
	LC/MS	=	liquid chromatography/mass spectroscopy
25	M	=	molar
	min	=	minute
	mL	=	milliliter
	mmol	=	millimols
	NTCP	=	Na ⁺ taurocholate cotransporting polypeptide
30	PBS	=	phosphate buffered saline
	PPTS	=	pyridinium <i>p</i> -toluenesulfonate
	THF	=	tetrahydrofuran
	TFA	=	trifluoroacetic acid
	TMSOTf	=	trimethylsilyltrifluoromethanesulfonate
35	Trisyl	=	2,4,6-triisopropylbenzenesulfonyl
	μL	=	microliter

μM = micromolar
v/v = volume to volume

EXPERIMENTAL METHODS

106001-89242660
5 The following examples illustrate how the synthesis of GABA analog
conjugates could be conducted in order to prepare compounds of formula (I). It
is contemplated that other drugs may be used in place of a GABA analog with
similar synthetic schemes. The syntheses described below are illustrated in
Figure 4-8. While GABA is not included within the scope of this invention,
these examples are illustrative of how other drugs could be readily coupled to a
10 bile acid in an analogous manner.

EXAMPLE 1

Synthesis of Cholic Acid Gabapentin Dipeptides (9)

Cholic acid (6) (408 mg, 1 mmol) was dissolved in anhydrous THF (10
mL) and triethylamine (0.167 mL, 1.2 mmol) added slowly with stirring. The
15 solution was cooled to -5°C in an ice-salt bath, and ethyl chloroformate (0.12
mL, 1.2 mmol) added slowly, maintaining the temperature between -5 to 0°C .
After addition was complete, the cold mixture was stirred for an additional 15
minutes. A solution containing an amino acid (7) (1.75 mmol) in 2N NaOH (2
mL) was added and the mixture stirred for an additional 60 min at -5 to 0°C .
20 After removal of the THF *in vacuo*, saturated NaHCO_3 (15 mL) was added and
the aqueous mixture washed with EtOAc (3 x 10 mL), then the pH adjusted to 3-
4 with citric acid. The product was extracted into EtOAc (3 x 15 mL), and the
combined organic phase dried over MgSO_4 , and concentrated to dryness. The
residue was purified by flash chromatography on silica gel (5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$)
25 to give the pure cholic acid adduct (8). This compound (0.4 mmol) was
dissolved in anhydrous THF (10 mL) and triethylamine (0.44 mmol) added
slowly with stirring. The solution was cooled to -5°C in an ice-salt bath, and
ethyl chloroformate (44 μL , 0.44 mmol) added slowly, maintaining the

temperature between -5 to 0°C . After addition was complete, the cold mixture was stirred for an additional 15 minutes. A solution containing gabapentin (**2**) (166 mg, 0.8 mmol) in 2N NaOH (3 mL) was added and the mixture stirred for an additional 60 min at -5 to 0°C . After removal of the THF *in vacuo*, saturated NaHCO₃ (10 mL) was added and the aqueous mixture washed with EtOAc (3 x 10 mL), then the pH adjusted to 3-4 with citric acid. The product was extracted into EtOAc (3 x 20 mL), and the combined organic phases dried over MgSO₄, and concentrated to dryness. The residue was purified by flash chromatography on silica gel (10% MeOH/CH₂Cl₂) to give the pure cholic acid gabapentin dipeptide derivative. The corresponding sodium salt (**9**) was prepared in quantitative yield by addition of a methanol solution of the acid to water containing 0.5N NaOH (1 eq.) and evaporation to dryness on a lyophilizer. Compounds were characterized by electrospray mass spectrometry as reported below:

Cholyl-Gly-Gabapentin (**9a**): MS (ESI): m/z 617.50 ($\text{M}-\text{H}^-$), 619.51 ($\text{M}+\text{H}^+$).
¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 3.81 (s, 2H), 3.34 (s, 2H), 2.28 (s, 2H), 1.03 (d, 3H, $J=6.4\text{Hz}$), 0.91 (s, 3H), 0.70 (s, 3H).

Cholyl-Ala-Gabapentin (**9b**): MS (ESI): m/z 631.50 ($\text{M}-\text{H}^-$), 633.52 ($\text{M}+\text{H}^+$).
¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 4.29 (m, 1H), 3.34 (s, 2H), 2.28 (s, 2H), 1.34 (d, 2H, $J=6.8\text{Hz}$), 1.01 (d, 3H, $J=6.4\text{Hz}$), 0.91 (s, 3H), 0.71 (s, 3H).

Cholyl-Val-Gabapentin (**9c**): MS (ESI): m/z 659.55 ($\text{M}-\text{H}^-$), 661.55 ($\text{M}+\text{H}^+$).
¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 4.26 (m, 1H), 3.34 (s, 2H), 2.27 (s, 2H), 1.02 (d, 3H, $J=6.4\text{Hz}$), 0.97 (d, 6H, $J=6.4\text{Hz}$), 0.91 (s, 3H), 0.71 (s, 3H).

Cholyl-Leu-Gabapentin (**9d**): MS (ESI): m/z 673.43 ($M-H^-$), 675.45 ($M+H^+$).

1H NMR (CD_3OD , 400 MHz, characteristic resonances only): 4.34 (m, 1H), 3.34 (s, 2H), 2.27 (s, 2H), 1.02 (d, 3H, $J=6.4Hz$), 0.97 (d, 3H, $J=6.4Hz$), 0.92 (d, 3H, $J=6.4Hz$), 0.91 (s, 3H), 0.70 (s, 3H).

5 Cholyl-Norleu-Gabapentin (**9e**): MS (ESI): m/z 659.56 ($M-H^-$), 661.57 ($M+H^+$).

1H NMR (CD_3OD , 400 MHz, characteristic resonances only): 4.26 (m, 1H), 3.34 (s, 2H), 2.27 (s, 2H), 1.02 (d, 3H, $J=6.4Hz$), 0.91 (s, 3H), 0.71 (s, 3H).

Cholyl- t BuGly-Gabapentin (**9f**): MS (ESI): m/z 673.58 ($M-H^-$), 675.58 ($M+H^+$).

10 1H NMR (CD_3OD , 400 MHz, characteristic resonances only): 4.20 (s, 1H), 3.34 (s, 2H), 2.29 (s, 2H), 1.01 (d, 3H, $J=6.4Hz$), 0.98 (s, 9H), 0.91 (s, 3H), 0.70 (s, 3H).

Cholyl-Phe-Gabapentin (**9g**): MS (ESI): m/z 707.47 ($M-H^-$), 709.36 ($M+H^+$).

15 1H NMR (CD_3OD , 400 MHz, characteristic resonances only): 7.26 (m, 5H), 4.59 (m, 1H), 3.34 (s, 2H), 3.25-2.95 (m, 2H), 2.18 (d, 2H, $J=7.2Hz$), 0.98 (d, 3H, $J=6.4Hz$), 0.91 (s, 3H), 0.68 (s, 3H).

Cholyl-Tyr-Gabapentin (**9h**): MS (ESI): m/z 723.42 ($M-H^-$), 725.42 ($M+H^+$).

20 1H NMR (CD_3OD , 400 MHz, characteristic resonances only): 7.06 (d, 2H, $J=8.8Hz$), 6.69 (d, 2H, $J=8.8Hz$), 4.51 (dd, 1H, $J=6.8Hz$, $J=8.8Hz$), 3.34 (s, 2H), 3.16-2.78 (m, 2H), 2.16 (d, 2H, $J=7.2Hz$), 0.98 (d, 3H, $J=6.4Hz$), 0.91 (s, 3H), 0.68 (s, 3H).

Cholyl-Ser-Gabapentin (**9i**): MS (ESI): m/z 647.42 ($M-H^-$), 649.41 ($M+H^+$).

¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 4.37 (m, 1H), 3.78 (m, 2H), 3.34 (s, 2H), 2.15 (s, 2H), 1.03 (d, 3H, J=6.4Hz), 0.91 (s, 3H), 0.71 (s, 3H).

Cholyl-Asp-Gabapentin (**9j**): MS (ESI): m/z 647.45 (M-H⁻), 649.45 (M+H⁺).

5 ¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 4.71 (m, 1H), 3.34 (s, 2H), 2.87-2.65 (m, 2H), 2.28 (s, 2H), 1.02 (d, 3H, J=6.4Hz), 0.91 (s, 3H), 0.71 (s, 3H).

Cholyl-Glu-Gabapentin (**9k**): MS (ESI): m/z 688.50 (M-H⁻), 690.54 (M+H⁺).

10 ¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 4.35 (m, 1H), 3.34 (s, 2H), 2.38 (t, 2H, J=7Hz), 1.02 (d, 3H, J=6.4Hz), 0.91 (s, 3H), 0.71 (s, 3H).

Cholyl-Asn-Gabapentin (**9l**): MS (ESI): m/z 674.43 (M-H⁻), 676.44 (M+H⁺).

15 ¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 4.29 (m, 1H), 3.34 (s, 2H), 2.92-2.69 (m, 2H), 2.28 (s, 2H), 1.02 (d, 3H, J=6.4Hz), 0.91 (s, 3H), 0.70 (s, 3H).

Cholyl-Lys-Gabapentin (**9m**): MS (ESI): m/z 688.50 (M-H⁻), 690.54 (M+H⁺).

¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 4.29 (m, 1H), 3.34 (s, 2H), 2.28 (s, 2H), 1.02 (d, 3H, J=6.4Hz), 0.91 (s, 3H), 0.70 (s, 3H).

20 Cholyl-β-Ala-Gabapentin (**9n**): MS (ESI): m/z 631.45 (M-H⁻), 633.45 (M+H⁺).

¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 3.34 (s, 2H), 3.20 (t, 2H, J=8Hz), 2.29 (s, 2H), 2.26 (t, 2H, J=8Hz), 1.02 (d, 3H, J=6.4Hz), 0.91 (s, 3H), 0.70 (s, 3H).

Cholyl-Gaba-Gabapentin (**9o**): MS (ESI): m/z 645.56 (M-H⁻), 647.57 (M+H⁺).

¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 3.34 (s, 2H), 3.41 (t, 2H, J=6.8Hz), 2.44 (t, 2H, J=6.8Hz), 2.28 (s, 2H), 1.01 (d, 3H, J=6.4Hz), 0.91 (s, 3H), 0.71 (s, 3H).

5 Ursodeoxycholyl-Phe-Gabapentin (**9p**): MS (ESI) m/z 691.6 (M-H⁻), 693.5 (M+H⁺).

¹H NMR (CD₃OD, 400 MHz, characteristic resonances only): 7.26 (m, 5H), 4.60 (m, 1H), 3.34 (s, 2H), 3.25-2.95 (m, 2H), 2.08 (d, 2H, J=7Hz), 0.95 (s, 3H), 0.91 (d, 3H, J=6.4Hz), 0.67 (s, 3H).

10 Alternative Synthesis of Cholic Acid Gabapentin Dipeptides (**9**)

An N-Fmoc-protected amino acid (**10**) (1 mmol) is dissolved in anhydrous THF (10 mL) and triethylamine (0.167 mL, 1.2 mmol) added slowly with stirring. The solution is cooled to -5°C in an ice-salt bath, and ethyl chloroformate (0.12 mL, 1.2 mmol) added slowly, maintaining the temperature
15 between -5 to 0°C. After addition is complete, the cold mixture is stirred for an additional 15 minutes. A solution containing gabapentin (**2**) (1.75 mmol) in 2N NaOH (2 mL) is added and the mixture stirred for an additional 60 min at -5 to 0°C. After removal of the THF *in vacuo*, saturated NaHCO₃ (15 mL) is added and the aqueous mixture washed with EtOAc (3 x 10 mL), then the pH adjusted
20 to 3-4 with citric acid. The product is extracted into EtOAc (3 x 15 mL), and the combined organic phase dried over MgSO₄, and concentrated to dryness. The residue is purified by flash chromatography on silica gel (5% MeOH/CH₂Cl₂) to give the pure protected dipeptide adduct. This compound is dissolved in dry NMP (4 mL) containing 5% (v/v) DBU and the solution stirred for 30 min to
25 give the dipeptide (**11**). Cholic acid (**6**) (1 mmol) is dissolved in anhydrous THF (10 mL) and triethylamine (0.167 mL, 1.2 mmol) added slowly with stirring. The solution is cooled to -5°C in an ice-salt bath, and ethyl chloroformate (0.12

mL, 1.2 mmol) added slowly, maintaining the temperature between -5 to 0°C. After addition is complete, the cold mixture is stirred for an additional 15 minutes. The above solution of dipeptide (11) is added and the mixture stirred for an additional 60 min at -5 to 0°C. After removal of the solvent *in vacuo*, saturated NaHCO₃ (15 mL) is added and the aqueous mixture washed with EtOAc (3 x 10 mL), then the pH adjusted to 3-4 with citric acid. The product is extracted into EtOAc (3 x 15 mL), and the combined organic phase dried over MgSO₄, and concentrated to dryness. The residue is purified by flash chromatography on silica gel (10% MeOH/CH₂Cl₂) to give the pure cholic acid gabapentin dipeptide derivative. The corresponding sodium salt (9) is prepared in quantitative yield by addition of a methanol solution of the acid to water containing 0.5N NaOH (1 eq.) and evaporation to dryness on a lyophilizer.

EXAMPLE 2

Synthesis of Cholic Acid Pregabalin Dipeptides (12)

Pregabalin (3), prepared according the methods described in Silverman *et al* (US Patent 5,563,175), is transformed to the cholyl dipeptide derivative (12) following the procedure detailed above for the gabapentin analog (9).

EXAMPLE 3

In Vitro Compound Transport Assays with

IBAT and LBAT-Expressing Cell Lines

(a) Inhibition of Radiolabeled Taurocholate Uptake

CHO cells transfected with the IBAT or LBAT transporter were seeded into 96-well microtiter plates at 100,000 cells/well in 100 µL DMEM containing 10% serum, glutamine and Penstrep. After overnight incubation the media was removed and test compound (25 µL) added at 2x the final desired concentration. Tritiated taurocholate (50,000 CPM/well) was diluted with cold substrate to a final concentration of 5 µM and 25 µL/well of this mixture was added to the

plate. After incubating for 1 h at room temperature the solution was removed and the plate washed 4x with PBS at 4°C. 200 µL/well of scintillant is added and the plate then read in a Wallac microbeta counter. The inhibition data is processed by standard methods to calculate an inhibition constant K_i for the test compound.

(b) Analysis of Electrogenic Transport in *Xenopus* Oocytes

RNA preparation: Human IBAT Transporter cDNAs were subcloned into a modified pGEM plasmid that contains 5' and 3' untranslated sequences from the *Xenopus* β -actin gene. These sequences increase RNA stability and protein expression. Plasmid cDNA was linearized and used as template for *in vitro* transcription (Epicentre Technologies transcription kit, 4:1 methylated:non-methylated GTP).

Xenopus oocyte isolation. *Xenopus laevis* frogs were anesthetized by immersion in Tricaine (1.5 g/mL in deionized water) for 15 min. Oocytes were removed and digested in frog ringer solution (90 mM NaCl, 2 mM KCl, 1 mM $MgCl_2$, 10 mM NaHEPES, pH 7.45, no $CaCl_2$) with 1 mg/mL collagenase (Worthington Type 3) for 80-100 min with shaking. The oocytes were washed 6 times, and the buffer changed to frog ringer solution containing $CaCl_2$ (1.8 mM). Remaining follicle cells were removed if necessary. Cells were incubated at 16°C, and each oocyte injected with 10-20 µg RNA in 45 µL solution.

Electrophysiology measurements. Transport currents were measured 2-14 days after injection, using a standard two-electrode electrophysiology set-up (Geneclamp 500 amplifier, Digidata 1320/PCLAMP software and ADInstruments hardware and software were used for signal acquisition). Electrodes (2-4 m Ω) were microfabricated using a Sutter Instrument puller and filled with 3M KCl. The bath was directly grounded (transporter currents were

less than 0.3 μ A). Bath flow was controlled by an automated perfusion system (ALA Scientific Instruments, solenoid valves).

For transporter pharmacology, oocytes were clamped at -60 to -90 mV, and continuous current measurements acquired using PowerLab Software and an ADInstruments digitizer. Current signals were lowpass filtered at 20 Hz and acquired at 4-8 Hz. All bath and drug-containing solutions were frog ringers solution containing CaCl_2 . Drugs were applied for 10-30 seconds until the induced current reached a new steady-state level, followed by a control solution until baseline currents returned to levels that preceded drug application. The difference current (baseline subtracted from peak current during drug application) reflected the net movement of charge resulting from electrogenic transport and was directly proportional to transport rate. Recordings were made from a single oocyte for up to 60 min, enabling 30-40 separate compounds to be tested per oocyte. Compound-induced currents were saturable and gave half-maximal values at substrate concentrations comparable to radiolabel competition experiments. To compare results between oocytes expressing different levels of transport activity, a saturating concentration of glycodeoxycholate (300 μ M) was used as a common reference to normalize results from test compounds. Using this normalization procedure V_{max} (i.e. maximal induced current) for different compounds tested on different oocytes could be compared.

Table 1: *In vitro* transport data for selected compounds on IBAT-expressing cells

COMPOUND	IC ₅₀ (μM)	EC ₅₀ (μM)	% Max. (GDC)
Cholyl-Gly-Gabapentin (9a)	66	22	67
Cholyl-Phe-Gabapentin (9g)	92	140	28
Cholyl-Tyr-Gabapentin (9h)	52	160	13
UDC-Phe-Gabapentin (9p)	27	18	27

IC₅₀ data from radiolabeled competition assay in transporter-expressing CHO cells

EC₅₀ and %Max data (relative to glycodeoxycholate) from transporter-expressing oocytes

Table 2: *In vitro* transport data for selected compounds on LBAT-expressing cells

COMPOUND	IC ₅₀ (μM)	EC ₅₀ (μM)	% Max. (GDC)
Cholyl-Gly-Gabapentin (9a)	64	ND	ND
Cholyl-Phe-Gabapentin (9g)	0.5	ND	ND
Cholyl-Tyr-Gabapentin (9h)	15	ND	ND
UDC-Phe-Gabapentin (9p)	0.7	ND	ND

IC₅₀ data from radiolabeled competition assay in transporter-expressing CHO cells

EC₅₀ and %Max data (relative to glycodeoxycholate) from transporter-expressing oocytes

EXAMPLE 4

In Vitro Compound Transport Assays with PEPT1 and PEPT2-Expressing Cell Lines - Inhibition of Radiolabeled Gly-Sar Uptake

Rat and human PEPT1 and PEPT2 expressing CHO cell lines were prepared as described in PCT Application WO01/20331. Gabapentin-containing dipeptides were evaluated for interaction with the peptide transporters using a radiolabeled substrate uptake assay in a competitive inhibition format, as described in PCT Application WO01/20331. Transport-induced currents were also measured in *Xenopus* oocytes transfected with rat and human PEPT1 and PEPT2.

Table 3: *In vitro* transport data for selected compounds on rPEPT1-expressing cells

COMPOUND	IC ₅₀ (μM)	% Max. (Gly-Sar)
Gly-Gabapentin	320	76
Phe-Gabapentin	56	52
Tyr-Gabapentin	130	22

IC₅₀ data from radiolabeled competition assay in transporter-expressing CHO cells

%Max response (relative to Gly-Sar) from transporter-expressing oocytes at a test compound concentration of 1 mM

Table 4: *In vitro* transport data for selected compounds on rPEPT2-expressing cells

COMPOUND	IC ₅₀ (μM)	% Max. (Gly-Sar)
Gly-Gabapentin	ND	ND
Phe-Gabapentin	ND	77
Tyr-Gabapentin	4	73

IC₅₀ data from radiolabeled competition assay in transporter-expressing CHO cells

%Max response (relative to Gly-Sar) from transporter-expressing oocytes at a test compound concentration of 1 mM

EXAMPLE 5

In Vitro Enzymatic Release of Gabapentin from Cholyl-Amino Acid-Gabapentin Conjugates

Sustained oral delivery of a drug molecule by attachment through a cleavable linker arm to an actively transported promoiety requires that the drug eventually be released from the drug/cleavable linker/transporter compound (prodrug) by enzymatic cleavage in one or more tissues of the enterohepatic circulation. The release of Gabapentin from the prodrug Cholyl-Phe-Gabapentin (**9g**) (and other Cholyl-Amino Acid-Gabapentin conjugates (**9**)) was evaluated *in vitro* using tissues representative of those involved in the enterohepatic

circulation. These studies indicated that *in vitro* cleavage of the prodrug could occur via a stepwise process, with release of the Gabapentin-containing dipeptide (e.g. Phe-Gabapentin) preceding hydrolysis to liberate free Gabapentin.

5 Tissues were obtained from commercial sources (e.g., Pel-Freez Biologicals, Rogers, AR, or GenTest Corporation, Woburn, MA). Stability of Choly-Phe-Gabapentin towards specific enzymes (e.g., carboxypeptidase A, cholyglycine hydrolase) was also evaluated by incubation with the purified enzyme.

10 Experimental conditions used for the *in vitro* studies are described in Table 3 below. Each preparation was incubated with (9g) at 37°C for one hour. Aliquots (50 µL) were removed at 0, 30, and 60 min and quenched with 0.1% trifluoroacetic acid in acetonitrile. Samples were then centrifuged and analyzed by LC/MS/MS as described below.

15 The stability of Gabapentin-containing dipeptides to purified aminopeptidase 1 and to Caco-2 homogenates was evaluated as follows:
Aminopeptidase Stability: Aminopeptidase 1 (Sigma catalog # A-9934) was diluted in deionised water to a concentration of 856 units/mL. Stability studies were conducted by incubating prodrug (5 µM) with 0.856 units/mL
20 aminopeptidase 1 in 50 mM Tris-HCl buffer at pH 8.0 and 37°C. Concentrations of intact prodrug and released drug were determined at zero time and 60 minutes using LC/MS/MS.

Pancreatin Stability: Stability studies were conducted by incubating prodrug (5 µM) with 1 % (w/v) pancreatin (Sigma, P-1625, from porcine
25 pancreas) in 0.025 M Tris buffer containing 0.5 M NaCl (pH 7.5) at 37 °C for 60

106001-89242660

min. The reaction was stopped by addition of 2 volumes of methanol. After centrifugation at 14,000 rpm for 10 min, the supernatant was removed and analyzed by LC/MS/MS.

5 Caco-2 Homogenate S9 Stability: Caco-2 cells were grown for 21 days
prior to harvesting. Culture medium was removed and cell monolayers were
rinsed and scraped off into ice-cold 10 mM sodium phosphate/0.15 M potassium
chloride, pH 7.4. Cells were lysed by sonication at 4°C using a probe sonicator.
Lysed cells were then transferred into 1.5 mL centrifuge vials and centrifuged at
9000 g for 20 min at 4°C. The resulting supernatant (Caco-2 cell homogenate S9
10 fraction) was aliquoted into 0.5 mL vials and stored at -80°C until used.
For stability studies, prodrug (5 µM) was incubated in Caco-2 homogenate S9
fraction (0.5 mg protein per mL) for 60 min at 37°C. Concentrations of intact
prodrug and released drug were determined at zero time and 60 minutes using
LC/MS/MS.

15 Concentrations of Cholyl-Phe-Gabapentin (**9g**), Phe-Gabapentin or
Gabapentin in tissue extracts were determined by direct injection onto an API
2000 LC/MS/MS equipped with an Agilent 1100 binary pump and autosampler.
Separation was achieved using a 3.5 µm Zorbax Ellipse XDB-C8 4.4 x 150 mm
column heated to 45°C during the analysis. The mobile phases were: 0.1%
20 formic acid in water (A) and 0.1% formic acid in acetonitrile (B). The gradient
condition was: 2% B for 0.5 min, increasing to 90% B in 2.0 min, maintained for
2.5 min and returning to 2% B for 2 min. A TurbolonSpray source was used on
the API 2000. The analysis was performed in the positive ion mode and MRM
transitions of 709.5/172.1 and 172.0/137.2 were used in the analysis of Cholyl-
25 Phe-Gabapentin (**9g**) and Gabapentin (**2**) respectively. Ten microliters of the
sample extracts were injected. Peaks were integrated using Analyst quantitation

software. The method was linear for (9g) or (2) over the concentration range 0.01 to 12.5 µg/mL and 0.002 to 2.5 µg/mL respectively.

Table 5. *In Vitro* Enzymatic Release of Phe-Gabapentin from Cholyl-Phe-Gabapentin (9g)

Preparation	Substrate Concentration	Cofactors	Percent of Phe-Gabapentin Released in 60 min
Rat Plasma	2.0 µM	None	NR
Human Plasma	2.0 µM	None	NR
Rat Liver S9 (0.5 mg/mL)	2.0 µM	NADPH	NR
Human Liver S9 (0.5 mg/mL)	2.0 µM	NADPH	NR
Human Intestine S9 (0.5 mg/mL)	2.0 µM	NADPH	NR
Cholylglycine Hydrolase (87 units/mL)	0.8 µM	None	~3
Carboxypeptidase A (10 units/mL)	2.0 µM	None	NR

NR = Not released

Table 6. *In Vitro* Enzymatic Release of Gabapentin (2) from Phe-Gabapentin

Preparation	Substrate Concentration	Cofactors	Percent of Gabapentin Released in 60 min
Rat Plasma	2.0 µM	None	19

Preparation	Substrate Concentration	Cofactors	Percent of Gabapentin Released in 60 min
Human Plasma	2.0 μ M	None	NR
Rat Liver S9 (0.5 mg/mL)	2.0 μ M	NADPH	1
Human Liver S9 (0.5 mg/mL)	2.0 μ M	NADPH	1
Human Intestine S9 (0.5 mg/mL)	2.0 μ M	NADPH	5
Cholylglycine Hydrolase (87 units/mL)	0.8 μ M	None	NR
Carboxypeptidase A (10 units/mL)	2.0 μ M	None	NR
Caco-2 Homogenate	5.0 μ M	None	21
Aminopeptidase	5.0 μ M	None	24

NR = Not released

Table 7: *In Vitro* Enzymatic Release of Gabapentin (2) from Cholyl-Amino Acid-Gabapentin Compounds by Pancreatin

COMPOUND	% (2) Released
Cholyl-Gly-Gabapentin (9a)	NR
Cholyl-Phe-Gabapentin (9g)	4
Cholyl-Tyr-Gabapentin (9h)	40

NR = Not released

EXAMPLE 6

Sustained Release of Gabapentin (2) from Cholyl-Phe-Gabapentin (9g) .

Following Oral Administration to Rats

The pharmacokinetics of the prodrug Cholyl-Phe-Gabapentin (9g) were examined in rats. Three groups of four male Sprague-Dawley rats (approx 200 g) with jugular cannulae each received one of the following treatments: A) a single bolus intravenous injection of Gabapentin (25 mg/kg, as a solution in

water); B) a single oral dose of Gabapentin (25 mg/kg, as a solution in water) administered by oral gavage; C) a single oral dose of (9g) (103.5 mg/kg, as a solution in water) administered by oral gavage. Animals were fasted overnight prior to dosing and until 4 hours post-dosing. Serial blood samples were obtained over 24 hours following dosing and blood was processed for plasma by centrifugation. Plasma samples were stored at -80°C until analyzed.

Concentrations of (9g) or (2) in plasma samples were determined by LC/MS/MS as described above. Plasma (50 μL) was precipitated by addition of 100 mL of methanol and supernatant was injected directly onto the LC/MS/MS system. Following oral administration of Gabapentin, concentrations of Gabapentin in plasma reached a maximum at 1.3 ± 0.5 hours (T_{max}) and declined thereafter with a terminal half-life of 3.0 ± 1.3 hours. The oral bioavailability of Gabapentin was $87 \pm 18\%$. Following oral administration of Choly-Phe-Gabapentin (9g), concentrations of Gabapentin in plasma reached a maximum at ~ 7.1 hours post-dosing and declined thereafter with a terminal half-life of ~ 5.1 hours. Concentrations of released Gabapentin in plasma were sustained beyond 24 hours. These data indicate that prodrug Choly-Phe-Gabapentin (9g) is metabolized to Gabapentin (2) *in vivo*, and that a substantially sustained release of Gabapentin was achieved following oral administration of (9g) compared to the relatively rapid clearance observed for oral Gabapentin.

EXAMPLE 7

Sustained Release of Gabapentin (2) from Choly-Gly-Gabapentin (9a) Following Oral Administration to Rats

Following oral administration of Choly-Gly-Gabapentin (9a) (90.4 mg/kg, as a solution in water) according to the protocol of Example 6 above, concentrations of Gabapentin in plasma reached a maximum at ~ 8.0 hours post-dosing and declined thereafter with a terminal half-life of ~ 6.9 hours.

Concentrations of released Gabapentin in plasma were sustained beyond 24 hours. These data indicate that prodrug Cholyl-Gly-Gabapentin (9a) is metabolized to Gabapentin (2) *in vivo*, and that a substantially sustained release of Gabapentin was achieved following oral administration of (9a) compared to the relatively rapid clearance observed for oral Gabapentin.

EXAMPLE 8

Sustained Release of Gabapentin (2) from Cholyl-Tyr-Gabapentin (9h) Following Oral Administration to Rats

Following oral administration of Cholyl-Tyr-Gabapentin (9h) (106 mg/kg, as a solution in water) according to the protocol of Example 6 above, concentrations of Gabapentin in plasma reached a maximum at ~ 7.3 hours post-dosing and declined thereafter with a terminal half-life of ~ 4.5 hours.

Concentrations of released Gabapentin in plasma were sustained beyond 24 hours. These data indicate that prodrug Cholyl-Tyr-Gabapentin (9h) is metabolized to Gabapentin (2) *in vivo*, and that a substantially sustained release of Gabapentin was achieved following oral administration of (9h) compared to the relatively rapid clearance observed for oral Gabapentin.

In view of the above disclosure, it is understood, of course, that combinations of substituents within the compounds of the present invention do not include any combination that is chemically impossible or non-feasible as would be appreciated by one skilled in the art.

REFERENCES

Arya, P.; Burton, G. W. Bile acids for biological and chemical applications and processes for the production thereof. U.S. Patent 5,541,348, 1996.

Baringhaus, K.-H.; Matter, H.; Stengelin, S.; Kramer, W. Substrate specificity of the ileal and hepatic Na⁺ / bile acid cotransporters of the rabbit. II. A reliable 3D QSAR pharmacophore model for the ileal Na⁺ / bile acid cotransporter. *J. Lipid Res.* **1999**, *40*, 2158-2168.

- 5 Bryans, J. S.; Wustrow, D. J. 3-Substituted GABA analogs with central nervous system activity: a review. *Med. Res. Rev.* **1999**, *19*, 149-177.

Bundgaard, H. in *Design of Prodrugs* (Bundgaard, H. Ed.), Elsevier Science B.V., **1985**, pp. 1-92.

- 10 Ho, N. F. H. Utilizing bile acid carrier mechanisms to enhance liver and small intestine absorption. *Ann. N. Y. Acad. Sci.* **1987**, *507*, 315-329.

Jezyk, N.; Li, C.; Stewart, B. H.; Wu, X.; Bockbrader, H. N.; Fleisher, D. Transport of pregabalin in rat intestine and Caco-2 monolayers. *Pharm. Res.* **1999**, *16*, 519-526.

- 15 Kagedahl, M.; Swaan, P. W.; Redemann, C. T.; Tang, M.; Craik, C. S.; Szoka, F. C.; Oie, S. Use of the intestinal bile acid transporter for the uptake of cholic acid conjugates with HIV-1 protease inhibitory activity. *Pharm. Res.* **1997**, *14*, 176-180.

- 20 Kim, D.-C.; Harrison, A. W.; Ruwart, M. J.; Wilkinson, K. F.; Fisher, J. F.; Hidalgo, I. J.; Borchardt, R. T. Evaluation of bile acid transporter in enhancing intestinal permeability of renin-inhibitory peptides. *J. Drug Targeting* **1993**, *1*, 347-359.

09374769-10001-89242660

Kramer, W.; Wess, G.; Schubert, G.; Bickel, M.; Girbig, F.; Gutjahr, U.; Kowalewski, S.; Baringhaus, K.-H.; Enhnen, A.; Glombik, H.; Mullner, S.; Neckermann, G.; Schulz, S.; Petzinger, E. Liver-specific drug targeting by coupling to bile acids. *J. Biol. Chem.* **1992**, *267*, 18598-18604.

- 5 Kramer, W.; Wess, G.; Neckermann, G.; Schubert, G.; Fink, J.; Girbig, F.; Gutjahr, U.; Kowalewski, S.; Baringhaus, K.-H.; Boger, G.; Enhnen, A.; Falk, E.; Friedrich, M.; Glombik, H.; Hoffmann, A.; Pittius, C.; Urmann, M. Intestinal absorption of peptides by coupling to bile acids. *J. Biol. Chem.* **1994a**, *269*, 10621-10627.

- 10 Kramer, W.; Wess, G.; Enhnen, A.; Bock, K.; Falk, E.; Hoffmann, A.; Neckerman, G.; Gantz, D.; Schulz, S.; Nickau, L.; Petzinger, E.; Turley, S.; Dietschy, J. M. Bile acid derived HMG-CoA reductase inhibitors. *Biochim. Biophys. Acta* **1994b**, *1227*, 137-154.

- 15 Kramer, W.; Wess, G. Modified bile acid conjugates, and their use as pharmaceuticals. U.S. Patent 5,462, 933, **1995**.

Kramer, W.; Wess, G. Bile acid conjugates of proline hydroxylase inhibitors. U.S. Patent 5,646,272, **1997a**.

Kramer, W.; Wess, G. Bile acid derivatives, processes for their preparation, and use as pharmaceuticals. U.S. Patent 5,668,126, **1997b**.

- 20 Kramer, W.; Stengelin, S.; Baringhaus, K.-H.; Enhnen, A.; Heuer, H.; Becker, W.; Corsiero, D.; Girbig, F.; Noll, R.; Weyland, C. Substrate specificity of the ileal and hepatic Na⁺ / bile acid cotransporters of the rabbit. I. Transport studies

with membrane vesicles and cell lines expressing the cloned transporters. *J. Lipid Res.* **1999**, *40*, 1604-1617.

Kullak-Ublick, G. A.; Beuers, U.; Paumgartner, G. Hepatobiliary transport. *J. Hepatology* **2000**, *32 (Suppl. 1)*, 3-18.

- 5 Mills, C. O; Iqbal, S.; Elias, E. Ileal absorption of tyrosine-conjugated bile acids in Wistar rats. *Biochim. Biophys. Acta* **1987**, *926*, 154-159.

Navia, M. A.; Chaturvedi, P. R. Design principles for orally bioavailable drugs. *Drug Discovery Today* **1996**, *1*, 179-189.

- 10 Petzinger, E.; Nickau, L.; Horz, J. A.; Schulz, S.; Wess, G.; Enhnen, A.; Falk, E.; Baringhaus, K.-H.; Glombik, H.; Hoffmann, A.; Mullner, S.; Neckermann, G.; Kramer, W. Hepatobiliary transport of hepatic 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors conjugated with bile acids. *Hepatology* **1995**, *22*, 1801-1811.

- 15 Reiner, A. Process for preparing ursodeoxycholic acid derivatives and their inorganic and organic salts having therapeutic activity. Eur. Patent 0 272 462 B1, **1992**.

Swaan, P. W.; Szoka, F. C.; Oie, S. Use of the intestinal and hepatic bile acid transporters for drug delivery. *Adv. Drug Delivery Rev.* **1996**, *20*, 59-82.

- 20 Swaan, P. W.; Hillgren, K. M.; Szoka, F. C.; Oie, S. Enhanced transepithelial transport of peptides by conjugation to cholic acid. *Bioconj. Chem.* **1997**, *8*, 520-525.

T06001-139242660

Tsuji, A.; Tamai, I. Carrier-mediated intestinal transport of drugs. *Pharm. Res.* **1996**, *13*, 963-977.

Each of the above references is incorporated herein by reference in its entirety.

094742660